

12/31/01

JC525  
S. PTOFORM PTO-1390  
(REV. 9-2001)

U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE

JC07 Rec'd PCT/PTO 31 DEC 2001 box 5eq

TRANSMITTAL LETTER TO THE UNITED STATES  
DESIGNATED/ELECTED OFFICE (DO/EO/US)  
CONCERNING A FILING UNDER 35 U.S.C. 371

ATTORNEY'S DOCKET NUMBER

522-1769

U.S. APPLICATION NO. (If known, see 37 CFR 1.5)

10/030522

INTERNATIONAL APPLICATION NO.

PCT/EP00/06677

INTERNATIONAL FILING DATE

13 JULY 2000

PRIORITY DATE CLAIMED

14 JULY 1999

TITLE OF INVENTION "Ligands for use in Therapeutic Compositions for the  
Treatment of Hemostasis Disorders"

APPLICANT(S) FOR DO/EO/US


JACQUEMIN, Marc G.; SAINT-REMY, Jean-Marie R.

Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:

1. ☒ This is a **FIRST** submission of items concerning a filing under 35 U.S.C. 371.
2. ☐ This is a **SECOND** or **SUBSEQUENT** submission of items concerning a filing under 35 U.S.C. 371.
3. ☐ This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.
4. ☒ The US has been elected by the expiration of 19 months from the priority date (Article 31).
5. ☒ A copy of the International Application as filed (35 U.S.C. 371(c)(2))
  - a. ☒ is attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ has been communicated by the International Bureau.
  - c. ☐ is not required, as the application was filed in the United States Receiving Office (RO/US).
6. ☐ An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).
  - a. ☐ is attached hereto.
  - b. ☐ has been previously submitted under 35 U.S.C. 154(d)(4).
7. ☒ Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))
  - a. ☒ are attached hereto (required only if not communicated by the International Bureau).
  - b. ☐ have been communicated by the International Bureau.
  - c. ☐ have not been made; however, the time limit for making such amendments has NOT expired.
  - d. ☐ have not been made and will not be made.
8. ☐ An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).
9. ☒ An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).
10. ☐ An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).

## Items 11 to 20 below concern document(s) or information included:

11. ☐ An Information Disclosure Statement under 37 CFR 1.97 and 1.98.
12. ☐ An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.
13. ☒ A **FIRST** preliminary amendment.
14. ☐ A **SECOND** or **SUBSEQUENT** preliminary amendment.
15. ☐ A substitute specification.
16. ☐ A change of power of attorney and/or address letter.
17. ☒ A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.
18. ☐ A second copy of the published international application under 35 U.S.C. 154(d)(4).
19. ☐ A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).
20. ☒ Other items or information:  
Priority Claim/Assertion of Small Entity Status

U.S. APPLICATION NO. <b>10/030522</b> INTERNATIONAL APPLICATION NO.			ATTORNEY'S DOCKET NUMBER			
21. <input checked="" type="checkbox"/> The following fees are submitted: <b>BASIC NATIONAL FEE (37 CFR 1.492 (a) (1) - (5)):</b> Neither international preliminary examination fee (37 CFR 1.482) nor international search fee (37 CFR 1.445(a)(2)) paid to USPTO and International Search Report not prepared by the EPO or JPO ..... <b>\$1040.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but International Search Report prepared by the EPO or JPO ..... <b>\$890.00</b>  International preliminary examination fee (37 CFR 1.482) not paid to USPTO but international search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... <b>\$740.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO but all claims did not satisfy provisions of PCT Article 33(1)-(4) ..... <b>\$710.00</b>  International preliminary examination fee (37 CFR 1.482) paid to USPTO and all claims satisfied provisions of PCT Article 33(1)-(4) ..... <b>\$100.00</b> <b>ENTER APPROPRIATE BASIC FEE AMOUNT =</b>			<b>CALCULATIONS PTO USE ONLY</b>          <table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width: 50%; text-align: right;">\$ 890.00</td> <td style="width: 50%;"></td> </tr> </table>		\$ 890.00	
\$ 890.00						
Surcharge of <b>\$130.00</b> for furnishing the oath or declaration later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(e)).			<table border="1" style="width:100%; border-collapse: collapse;"> <tr> <td style="width: 50%; text-align: right;">\$</td> <td style="width: 50%;"></td> </tr> </table>		\$	
\$						
CLAIMS	NUMBER FILED	NUMBER EXTRA	RATE	\$		
Total claims	22 - 20 =	2	x \$18.00	\$36.00		
Independent claims	9 - 3 =	6	x \$84.00	\$504.00		
MULTIPLE DEPENDENT CLAIM(S) (if applicable)			+ \$280.00	\$		
<b>TOTAL OF ABOVE CALCULATIONS =</b>				\$1430		
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27. The fees indicated above are reduced by 1/2.			+	\$		
<b>SUBTOTAL =</b>				\$		
Processing fee of <b>\$130.00</b> for furnishing the English translation later than <input type="checkbox"/> 20 <input type="checkbox"/> 30 months from the earliest claimed priority date (37 CFR 1.492(f)).			\$			
<b>TOTAL NATIONAL FEE =</b>				\$715		
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property			+	\$		
<b>TOTAL FEES ENCLOSED =</b>				\$715		
				Amount to be refunded:	\$	
				charged:	\$	
a. <input checked="" type="checkbox"/> A check in the amount of \$ <u>715.00</u> to cover the above fees is enclosed.  b. <input type="checkbox"/> Please charge my Deposit Account No. _____ in the amount of \$ _____ to cover the above fees. A duplicate copy of this sheet is enclosed.  c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>12-0913</u> . A duplicate copy of this sheet is enclosed.  d. <input type="checkbox"/> Fees are to be charged to a credit card. <b>WARNING:</b> Information on this form may become public. <b>Credit card</b> <b>information should not be included on this form.</b> Provide credit card information and authorization on PTO-2038.						
NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137 (a) or (b)) must be filed and granted to restore the application to pending status.  SEND ALL CORRESPONDENCE TO:						
				 SIGNATURE <u>MARK A. HAGEDORN</u> NAME <u>44,731</u> REGISTRATION NUMBER		

Please acknowledge  
receipt of the  
enclosed:

JACQUEMIN, Marc G.

SAINT-REMY, Jean-Marie R.

SERIAL NO: To Be Assigned

FILE DATE: Herewith

1. Express Mail Cert # EL 874 025 580 US
2. Transmittal Letter to the U.S. Designated  
Office (DO/EO/US) Form PTO-1390
3. Return Post Card
4. Amendment Accompanying Application
5. Priority Claim
6. Check No. 47420 in the amount of \$ 715.00
7. International Preliminary Examination Report  
(PCT/EPOO/06677)
8. Corrected Sequence Listing (paper copy, pages  
1-9)
9. Original Sequence Listing in Computer  
Readable Form (CRF) (3.5" diskettes (2))
10. Corrected Sequence listing in CRF (3.5"  
Diskettes (2))
11. Application (PCT/EPOO/06677)

DUE DATE (If Any):

DATE SENT:

December 31, 2001

MAH/td

"Express Mail" mailing label number


EL 874 025 580 US

Date of deposit: December 31, 2001

I hereby certify that this paper or fee is being deposited with  
the United States Postal Service "Express Mail Post Office  
to Addressee" service under 37 CFR 1.10 on the date indi-  
cated above and is addressed to the Assistant Commissioner  
for Patents, Washington DC 20231

Richard Sensenbrenner

(Typed or printed name of person mailing paper or fee)



(Signature of person mailing paper or fee)

10030522 050201

0522-1769

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

IN RE THE APPLICATION OF )

Jaquemin, Marc G. & )  
Saint-Remy, Jean-Marie R. )

Examiner:

Group Art Unit:

SERIAL NO.: To Be Assigned )

FILED: Herewith )

FOR: "Ligands for Use in Therapeutic )  
Compositions for the Treatment of )  
Hemostasis Disorders" )

**CERTIFICATE OF MAILING**

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to "Director of Patents and Trademarks, Washington, D.C. 20231," on December 31, 2001.

Name of person signing Terr Dickinson

Signature 

Honorable Director of  
Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

**PRELIMINARY AMENDMENT ACCOMPANYING APPLICATION**

Dear Sir:

The present application is the national filing of International Application No. PCT/EP00/06677 filed 13 July 2000. Appended hereto is a copy of the International Preliminary Examination Report for that application which is amended thereto the claims as they presently appear in the application. Before calculation of the national filing fee for the United States, it is requested that the application be amended as follows:

### **In the Claims**

Please cancels claims 1-26 and please add new claims 27-48 as follows:

27. A cell line named KRIX 1 deposited with the Belgian Coordinated Collections of Micro-organisms, under accession number LMBP 5089CB.
28. A monoclonal antibody having the capacity of at least 65% and at most 98% inactivating factor VIII or a complex of two or more factors involving factor VIII when the said monoclonal antibody is in a physiological excess by binding to a site of the said factor or complex.
29. A cell line producing a human monoclonal antibody having the capacity of at least 65% and at most 98% inactivating factor VIII or a complex of two or more factors involving factor VIII when the said monoclonal antibody is in a physiological excess by binding to a site of the said factor or complex.
30. A monoclonal antibody according to claim 28, wherein the binding site of the monoclonal antibody is not directly involved in a physiological interaction of the said factor or complex.
31. A monoclonal antibody according to claim 28, being able to recognize an epitope located in the C1 domain of factor VIII.
32. A monoclonal antibody according to claim 28, being produced by on purpose immunization in animals.
33. A monoclonal antibody according to claim 28, being produced by on purpose immunization in animals and being humanized.
34. An antigen-binding fragment Fab, Fab' or F(ab')<sub>2</sub>, a complementarity determining region, a soluble or membrane-anchored single-chain variable part, a single variable domain or a derivative of a monoclonal antibody having the capacity of at least 65% and at most 98% inactivating factor VIII or a complex of two or more factors involving factor VIII when

the said monoclonal antibody is in a physiological excess by binding to a site of the said factor or complex.

35. A pharmaceutical composition for the prevention or treatment of disorders of hemostasis and resulting pathologic conditions in mammals, comprising as an active ingredient a monoclonal antibody having the capacity of at least 65% and at most 98% inactivating factor VIII or a complex of two or more factors involving factor VIII when the said monoclonal antibody is in a physiological excess by binding to a site of the said factor or complex, or a fragment, derivative or homolog thereof, in admixture with a pharmaceutically acceptable carrier.
36. A pharmaceutical composition according to claim 35, further comprising a therapeutically effective amount of a thrombolytic agent.
37. A polynucleotide encoding for an antigen-binding fragment Fab, Fab' or F(ab')<sub>2</sub> or a complementarity determining region or a soluble or membrane-anchored single-chain variable part or a single variable domain or a derivative of a monoclonal antibody having the capacity of at least 65% and at most 98% inactivating factor VIII or a complex of two or more factors involving factor VIII when the said monoclonal antibody is in a physiological excess by binding to a site of the said factor or complex.
38. A method of obtaining monoclonal antibodies from a non-human mammal, comprising the steps of:
- a) selecting a non-human mammal having a modified and partially functional protein, the modification being with respect to a wild type protein and lying in a domain of the protein;
  - b) administering the wild type protein to the non-human mammal in order to elicit an immune response, and
  - c) selecting B-lymphocytes from the non-human mammal which produce antibodies which only partially inactivate the wild type protein.
39. A method of obtaining monoclonal antibodies from the blood of a human being having a modified and partially functional protein, the modification being with respect to a wild type protein and lying in a domain of the protein, and to whom the wild type protein was administered, the said method comprising the step of selecting, from the blood of said

human being, B-lymphocytes which produce antibodies which only partially inactivate the wild type protein.

40. A method according to claim 39, wherein the wild type protein is a factor or a complex of two or more factors involved in the coagulation cascade of blood.

41. A method according to claim 39, wherein the wild type protein is factor VIII or a complex including factor VIII.

42. A method according to claim 39, wherein the wild type protein is a protein involved in a proteolytic cascade.

43. A method according to claim 39, wherein the wild type protein is a complement factor.

44. A method of treatment and/or prevention of a disorder of hemostasis, coagulation disorder or thrombotic pathologic condition or attenuation of coagulation in a mammal, comprising administering to a mammal in need of such treatment or prevention or attenuation of coagulation a therapeutically effective amount of an active ingredient selected from a monoclonal antibody having the capacity of at least 65% and at most 98% inactivating factor VIII or a complex of two or more factors involving factor VIII when the said monoclonal antibody is in a physiological excess by binding to a site of the said factor or complex, or a fragment, derivative or homolog thereof.

45. A method according to claim 44, wherein the thrombotic pathologic condition is selected from intravascular coagulation, arterial thrombosis, arterial restenosis, venous thrombosis and arteriosclerosis.

46. A method according to claim 44, wherein the active ingredient is provided to the mammal by oral, intranasal, subcutaneous, intramuscular, intradermal, intravenous, intraarterial or parenteral administration or by catheterization.

47. A method according to claim 44, further comprising administering to the mammal, simultaneously or sequentially with the said active ingredient, a therapeutically effective amount of a thrombolytic agent.

48. A monoclonal antibody according to claim 28, having the capacity of inactivating the co-factor activity of factor VIII by targeting the C1 domain of factor VIII.

### **Sequence Listing**

Please cancel the sequence listing pages 1-7, as originally filed, and replace with the corrected replacement pages 1-9 appended hereto.

### **Remarks**

The above claim amendments are being made in order to eliminate multiple dependency and improper multiple dependency before calculation of the national filing fee for the United States. Should a multiple dependency remain, that is unintended, and the Patent and Trademark Office is requested to cancel any remaining multiple dependant claims without prejudice before calculation of the application filing fee. Applicants hereby assert small entity status.

Pursuant to 37 CFR § 1.825 amendment and correction of the sequence listing, as filed, is hereby made by replacement pages 1-9 attached in paper form, replacing pages 1-7 of the sequence listing contained in the application as filed. Applicants also enclose with the present application the original sequence listing, as filed in the international application, in Computer Readable Format (two copies on 3.5 inch blue diskettes, labeled as copy 1 and 2) and the corrected replacement sequence listing in Computer Readable Format (two copies on 3.5 inch green diskettes). The correction occurs on page 3 of the replacement sheets for SEQ ID No. 3.



The proper number of residues is 142, thus requiring the insertion of the additional residues after residue 60, thus becoming residues 61-80. Support for the amendment is found in Figure 7 of the specification, which properly includes the 142 residues. No new matter is included. The sequence listing information recorded in computer readable form is identical to the written sequence listing of the corresponding listing (original/replacement).

The International Preliminary Examination Report reaches the conclusion that claims meet the requirements of novelty and inventive step (nonobviousness). It is submitted that the same results should occur in the United States.

Examination of the applicant on its merits is awaiting.

Respectfully Submitted,

*December 31, 2001*

Date



Mark A. Hagedorn  
Registration No. 44,731  
LEE, MANN, SMITH, MCWILLIAMS,  
SWEENEY & OHLSON  
209 S. LaSalle Street, Suite 410  
Chicago, IL 60604-1202  
(312)368-1300  
(312)368-0064 (fax)

LIGANDS FOR USE IN THERAPEUTIC COMPOSITIONS FOR THE  
TREATMENT OF HEMOSTASIS DISORDERS.FIELD OF THE INVENTION

5 The present invention relates to novel cell lines and to ligands, namely human and/or humanized monoclonal antibodies, as well as fragments such as Fab, Fab', F(ab')<sub>2</sub>, scFv, single variable domains, complementarity determining regions, derivatives, homologs and combinations thereof, obtainable from the said cell lines. It also relates to pharmaceutical compositions comprising said ligands and to methods of  
10 preventing and treating coagulation disorders and resulting thrombotic pathologic conditions in humans by administration of the said ligands to patients in need thereof. It also relates to methods of obtaining specific mammalian antibodies.

BACKGROUND OF THE INVENTION

15 The formation of blood clots does not only limit bleeding in case of an injury (hemostasis), but may lead to serious organ damage and death in the context of atherosclerotic diseases by occlusion of an important artery or vein. Thrombosis is thus blood clot formation at the wrong time and place. It involves a cascade of complicated and regulated biochemical reactions between circulating blood proteins (coagulation  
20 factors), blood cells (in particular platelets) and elements of an injured vessel wall. Anticoagulation and antithrombotic treatment aim at inhibiting the formation of blood clots in order to prevent these dangerous consequences, such as myocardial infarction, stroke, loss of a limb by peripheral artery disease or pulmonary embolism. Given the importance of these diseases, it is rather surprising that antithrombotic therapy has  
25 relied on a few drugs since many years, namely Aspirin to inhibit platelets, Heparin that indirectly inhibits the coagulation factors IX, X and II (thrombin), and oral Warfarin that inhibits Vit K-dependent factors (VII, IX, X, II and Prot C). More recently, low molecular weight Heparins (inhibiting factors X and II to various degrees) have become anticoagulants of choice, largely because of their ease of application (once a day  
30 subcutaneous injection with no monitoring need). With growing understanding of the processes involved in thrombosis a growing number of specific inhibitors of coagulation factors have been developed. However, a better efficacy/safety ratio could to date not be obtained with them. Direct thrombin inhibitors, in particular, were linked to increased bleeding complications in large clinical trials.

35 Aspirin also provides a protective effect against thrombosis. It induces a long-lasting functional defect in platelets, detectable clinically as a prolongation of the bleeding time, through inhibition of the cyclooxygenase activity of the human platelet enzyme prostaglandin H-synthase (PGHS-1) with doses as low as 30 to 75 mg. Since

gastrointestinal side effects of aspirin appear to be dose-dependent, and for secondary prevention, treatment with aspirin is recommended for an indefinite period, there are practical reasons to choose the lowest effective dose. Further it has been speculated that a low dose (30 mg daily) might be more anti-thrombotic but attempts to identify the optimal dosage have yielded conflicting results. It has been claimed that the dose of aspirin needed to suppress fully platelet aggregation may be higher in patients with cerebrovascular disease than in healthy subjects and may vary from time to time in the same patient. However, aspirin in any daily dose of 30 mg or higher reduces the risk of major vascular events by 20 % at most, which leaves much room for improvement.

Further, the inhibiting role of aspirin may lead to prevention of thrombosis as well as to excess bleeding. The balance between the two depends critically on the absolute thrombotic versus hemorrhage risk of the patient.

In patients with acute myocardial infarction, reduction of infarct size, preservation of ventricular function and reduction in mortality has been demonstrated with various thrombolytic agents. However these agents still have significant shortcomings, including the need for large therapeutic doses, limited fibrin specificity, and significant associated bleeding tendency. Recombinant tissue plasminogen activator (t-PA) restores complete patency in just over one half of patients, whereas streptokinase achieves this goal in less than one third. Further, reocclusion after thrombolytic therapy occurs in 5 to 10 % of cases during the hospital stay and in up to 30 % within the first year according to Verheugt et al., *J. Am. Coll. Cardiol.* (1996) 27:618-627. Thus numerous studies have examined the effects of adjunctive antithrombin therapy for patients with acute myocardial infarction. As an example, U.S. Patent 5,589,173 discloses a method for dissolving and preventing reformation of an occluding thrombus comprising administering a tissue factor protein antagonist, which may be a monoclonal or polyclonal antibody, in adjunction to a thrombolytic agent.

Monoclonal antibodies have already been shown to be of therapeutic value as antithrombotic agents. The first approved drug in this field was Abciximab (ReoPro<sup>TM</sup>), a humanized Fab fragment of a murine monoclonal antibody (7E3) against platelet GP IIb/IIIa receptors. Murine antibodies have characteristics which may severely limit their use in human therapy. As foreign proteins, they may elicit an anti-immunoglobulin response termed human anti-mouse antibody (HAMA) that reduces or destroys their therapeutic efficacy and/or provokes allergic or hypersensitivity reactions in patients, as taught by Jaffers et al., *Transplantation* (1986) 41:572. The need for readministration in therapies of thromboembolic disorders increases the likelihood of such immune reactions. While the use of human monoclonal antibodies would address this limitation, it has proven difficult to generate large amounts of such antibodies by conventional hybridoma technology.

Recombinant technology has therefore been used to construct "humanized" antibodies that maintain the high binding affinity of murine monoclonal antibodies but exhibit reduced immunogenicity in humans. In particular, there have been suggested chimeric antibodies in which the variable region (V) of a non-human antibody is combined with the constant (C) region of a human antibody. As an example, the murine Fc fragment was removed from 7 E3 and replaced by the human constant immunoglobulin G Fab region to form a chimera known as c7 E3 Fab or abciximab. Methods of obtaining such chimerical immunoglobulins is described in detail in U.S. Patent 5,770,198.

The potential for synergism between GPIIb/IIIa inhibition by monoclonal antibody 7 E3 Fab and thrombolytic therapy was evaluated by Kleiman et al., *J. Am. Coll. Cardiol* (1993) 22:381-389. Major bleeding was frequent in this study. Hence, the potential for life-threatening bleeding is clearly a major concern with this combination of powerful antithrombotic compounds.

In a recent attempt to reduce the immunogenicity of murine antibodies, only the complementarity determining region (CDR), i.e. regions of hypervariability in the V regions, rather than the entire V domain, are transplanted to a human antibody. Such humanized antibodies are known as CDR-grafted antibodies. Such one CDR-grafted antibody was successfully constructed against the relatively simple nitrophenacetyl antigen, however the construction of CDR-grafted antibodies recognizing more complex antigens has resulted in antibodies having binding activity significantly lower than the native non-human antibodies. In numerous cases it was demonstrated that the mere introduction of non-human CDRs into a human antibody is insufficient to maintain full binding activity. While a refined computer model of the murine antibody of interest is required in order to identify critical amino-acids to be considered in the design of a humanized antibody, and general theoretical guidelines were proposed for such design, in all cases the procedure must be tailored and optimized for the particular non-human antibody of interest.

Tissue factor (TF), being a membrane glycoprotein functioning as a receptor for factor VII and VIIa and thereby initiating the said extrinsic pathway, has been investigated as a target for anticoagulant therapy. In addition to this role, TF has been implicated in pathogenic conditions such as vascular disease and gram-negative septic shock. A study attempting to characterize the anticoagulant potential of murine monoclonal antibodies showed that the inhibition of TF function by most of the monoclonal antibodies assessed was dependent upon the dissociation of the TF/VIIa complex that is rapidly formed when TF contacts plasma. One monoclonal antibody, TF8-5G9, was capable of inhibiting the TF/VIIa complex without dissociation of the

complex, thus providing an immediate anticoagulant effect in plasma, as disclosed in WO 96/40,921.

Targeted clotting factors exhibit both a medium molecular weight range (about 45,000 to 160,000) and a relatively high normal plasma concentration (at least 0.01 micromol/L).

One persistent concern with all available anti-thrombotic agents is the risk of overdose and therefore of excessive and life-threatening bleeding. Most current antithrombotic agents therefore warrant close monitoring of the patient.

Thus, there is a need for efficient compounds for the treatment of coagulation disorders, which cannot be overdosed, require no monitoring and are free from bleeding problems. For a therapeutic agent based on antibodies, the ideal compound would be a human antibody with full anticoagulant efficacy that does not induce immunogenicity.

Factor VIII is a protein providing important coagulant cofactor activity and is one of human clotting factors with a rather high molecular weight (265,000) and a very low normal plasma concentration (0.0007 micromol./litre). With its 2,332 amino-acid residues, factor VIII is one of the longest known polypeptide chains and is synthesized in the liver, the spleen and the placenta. Its gene has been shown to include 186,000 nucleotides.

Factor VIII circulates as inactive plasma protein. Factors V and VIII are homologous proteins sharing a common structural configuration of triplicated A domains and duplicated C domains with structurally divergent B domains connecting the A2 and A3 domains. Factor VIII circulates in a multiplicity of fragmented species in a tightly associated complex with von Willebrand factor at a concentration of 1 nmol/L. Factor VIII activation occurs by a cleavage between the A1 and A2 domains, resulting in the unstable heterotrimeric factor VIIIa molecule. Factor VIIIa binds tightly to membranes that contain acidic phospholipids. Factor VIII contains a phospholipid binding site in the C2 domain, between amino-acids 2302 and 2332, according to Arai et al. in *J.Clin.Invest.* (1989) 83:1978. Within the same factor VIII region, there is also a von Willebrand factor binding site acting in conjunction with amino-acid residues 1645-1689 in the A3 domain according to Shima et al. in *Throm.Haemost.* (1993) 69:240 and *J.Biol. Chem.* (1994) 269:11601.

Polyclonal antibodies inhibiting the co-factor activity of factor VIII have been classified as type I or type II inhibitors according to their capacity to inhibit factor VIII either completely (type I) or only partially (type II). According to Gawryl et al., *Blood* (1982) 60:1103-9, the reduced inactivation of factor VIII by human type II autoantibodies is believed to be due to a steric effect of von Willebrand factor. Monoclonal antibodies are not mentioned and, to date, no therapeutic use was made of such type II inhibitors. Biggs et al., *Br.J.Haematol.* (1972) 23:137 previously provided

an interpretation derived from data obtained by using human polyclonal antibodies, that a type II inhibitory pattern could be related to low affinity. B.Ly et al., *Scandinavian Journal of Haematology* (1982), 28:132-140 discloses polyclonal antibodies to factor VIII which most often belong to the IgG class both in hemophiliacs developing alloantibodies and in the more rare patients having autoantibodies against their own factor VIII. These polyclonal antibodies partially inactivate Factor VIII activity like the antibodies described in Biggs et al. (1972) and Hoyer et al. (1982). This document again fails to mention whether monoclonal antibodies can reproduce the pattern of Factor VIII inactivation shown by patient's polyclonal antibodies. Again, no monoclonal antibodies are mentioned-

European patent applications EP-A-123,945, EP-A-152,746 and EP-A-432,134 all disclose monoclonal antibodies produced by hybridoma cell lines and having a specific reactivity pattern with factor VIIIc polypeptide fragments. These monoclonal antibodies are said to be useful for detecting the presence of factor VIIIc and related polypeptides in plasma by immunoassay techniques, but a therapeutic potential use is not suggested in these documents.

J.Battle et al., *Annals of Hematology* (1997) 75:111-115, discloses a polyclonal alloantibody from a patient with severe von Willebrand disease showing, like a rabbit polyclonal antibody against von Willebrand factor, a partial inhibitory activity to plasma Factor VIII. These polyclonal anti-factor VIII antibodies therefore inactivate factor VIII following a pattern similar to anti-factor VIII type II antibodies found in patients with hemophilia A (Gawryl et al., *Blood* (1982) 60:1103-9). However, factor VIII antibodies were not detected in the said human alloantibody, thus suggesting that it was a non specific inhibition.

J.Ingerslev et al., *Clinica Chimica Acta* (1988) 174:65-82 discloses a series of murine monoclonal antibodies against human von Willebrand factor: two of them, belonging to the immunoglobulin isotype IgG1, exhibit an extremely low (1.3 BU/mg immunoglobulin) inhibition of factor VIII as shown in table I of said document. By comparison, human monoclonal antibody BO2C11, derived from a hemophilia A patient with inhibitor, has a specific activity of 7,000 BU/ mg protein (Jacquemin et al. *Blood*, (1998) 92:496-506). This indicates that administration of antibodies as described by Ingerslev to an animal or a human being would not affect factor VIII activity, unless an extremely high amount of antibody (hundreds of mg/ml) was present in plasma. The authors do not disclose whether when used in large excess these antibodies exhibit inhibitory activity like type I or type II (i.e. partial inactivation) polyclonal human factor VIII inhibitor, such as described in Gawryl et al., *Blood* (1982) 60:1103-9.

Maraganore et al., *Circulation* (1992) 86:413, showed that a synthetic 12-aminoacid peptide corresponding to residues 1675-1686 of factor VIII inhibits cleavage

by thrombin of the heavy chain required for the activation of the procoagulant activity of factor VIII and also of the light chain required to dissociate factor VIII from von Willebrand factor and that tyrosine sulfation of said peptide potentiates its recognition by factor VIII.

5        *J.Clin.Invest.* (1988) 82:206-211 describes obtaining an animal model for hemophilia A by infusion of human anti-factor VIII antibody in rabbits. According to WO 95/01570, antibodies against the light chain of human or porcine factor VIII were produced in a first animal and subsequently a temporary hemophilia was induced in a second animal by means of the purified monospecific antibody obtained. U.S. Patent  
10    5,804,159 also discloses inducing a temporary clotting disorder in a mammal by means of an anti-plasma antibody preparation acting on several blood coagulation factors, e.g. a preparation comprising antibodies against human von Willebrand factor and factor VIII, or against factor VIII/von Willebrand factor-complex, or against procoagulants, anticoagulants, clot structure factors, fibrinolysis factors and phospholipids.

15        However, none of the above-mentioned antibodies compounds involving factor VIII have been described for therapeutic purposes. In fact there is a prejudice among those skilled in the art against investigating anti-factor VIII antibodies for anti-thrombotic therapy because it is assumed that, a deficiency in factor VIII being the cause of hemophilia A, such antibodies would induce a bleeding state.

20        WO97/26010 discloses monoclonal antibodies having self-limiting neutralizing activity against a coagulation factor which are useful in pharmaceutical compositions for thrombotic disorders. Self-limiting neutralizing activity in this document is defined as the activity of an antibody that binds to a human coagulation factor and inhibits thrombosis in a manner such that limited modulation of coagulation is produced.  
25    Limited modulation of coagulation in turn is defined as an increase in clotting time as measured by prolongation of the activated partial thromboplastin time (aPTT) where plasma remains clottable with aPTT reaching a maximal value, preferably 35 to 100 seconds, despite increasing concentrations of the monoclonal antibody. APTT is thus used as the primary criterion for the evaluation of efficacy versus bleeding liability of  
30    antithrombotic agents.

More particularly, this document demonstrates that a sheep polyclonal to factor VIII (SAF8C-IG, purchased from Affinity Biologicals) induces a self-limiting prolongation of aPTT (the aPTT increased to a maximum of about 65 seconds). We have demonstrated, however, that SAF8C-IG totally inhibits the activity of human  
35    factor VIII (see Figure 10), i.e. is a type I inhibitor in the classification of Gawryl et al., Blood (1982) 60: 1103-9. This demonstrates that a limited increase in clotting time up to a certain maximum value is not necessarily correlated with partial inactivation of a clotting factor, and far less to a decrease in the risk of bleeding. For instance, it is well

known that patients with a complete deficit of coagulation factors have a limited prolongation of aPTT, usually in the area of 60 to 100 seconds, but are nevertheless exposed to a dramatic risk of bleeding (Hathaway et al. Am J Clin Pathol (1979) 71: 22-25, and Hoffmann et al. Thromb Haemostas (1978) 39: 640-645).

5           Conversely, it is well known that a prolonged APTT does not provide a valid parameter of the reduction of thrombosis risk. Notably, deficiency in factor XII, another coagulation factor of the intrinsic coagulation pathway results in APTT prolonged up to 6-fold (Hathaway et al. Am J Clin Pathol (1979) 71: 22-25, and; Hoffmann et al. Thromb Haemostas (1978) 39: 640-645). However, a significant number of patients  
10       with this deficiency have experienced myocardial infarction or thromboembolism, demonstrating the lack of protection from thrombotic disease in patient deficient in factor XII, despite important prolongation of the APTT (McPherson RA Am J Clin Pathol (1977) 68: 420, and; Glueck HI et al. Ann Intern Med (1966) 64:390).

          Jacquemin et al. in *Blood* (1998) 92:496-506 refers to a factor VIII-specific  
15       human IgG4 monoclonal antibody (BO2C11) produced by a cell line derived from the memory B-cell repertoire of a hemophilia A patient with inhibitors. BO2C11 is said to recognize the C2 domain of factor VIII and to inhibit its binding to both von Willebrand factor and phospholipids. It is said to completely inhibit the procoagulant activity of native and activated factor VIII with a specific activity of 7,000 Bethesda units/mg. The  
20       present inventors have further-shown that BO2C11, while totally inhibiting the activity of human factor VIII, provides a prolongation of about 110 seconds in clotting time as measured by aPTT, which again demonstrates that an increase in clotting time up to a certain maximum value is not necessarily correlated to partial inactivation of a coagulation factor. Such a reduction of factor VIII levels would expose the patient to  
25       severe risks of bleeding, like in patients with severe hemophilia A (Levine PH Ann NY Acad Sci (1975) 240:201; Gilbert MS Mount Sinai J Med (1977) 44: 339).

#### SUMMARY OF THE INVENTION

          The present invention is related to new ligands, namely new monoclonal human  
30       or humanized antibodies, fragments, derivatives and homologs thereof, which bind to a factor involved in hemostasis, in particular to a factor or factors of the coagulation cascade and more in particular bind to factor VIII or a complex thereof; to polypeptides or other molecules which bind to a factor or factors in hemostasis; to a novel cell line from which said monoclonal antibodies may be obtained; to pharmaceutical  
35       compositions comprising said ligands and to methods of prevention and treatment of coagulation disorders and resulting thrombotic pathologic conditions in humans by the administration of said ligands to patients in need thereof.



A first main object of the present invention is therefore to provide an effective and safe anti-thrombotic therapy which reduces the risk of bleeding in mammals, more particularly in humans.

It is a further object of this invention to provide therapeutic compositions which provide an effective anti-thrombotic therapy which reduces the risk of bleeding in mammals, more particularly in humans.

It is still a further object of the present invention to provide an anti-thrombotic therapy and anti-thrombotic therapeutic compounds which are safer to use than the previously known therapies and compositions.

One aspect of the present invention is to target a human protein factor involved in hemostasis, in particular in the coagulation cascade, more particularly factor VIII or a complex thereof, using specific ligands. Preferably, these ligands, being other than polyclonal antibodies, provide a therapeutically useful plateau level by only partially inhibiting the function of the targeted factor so that a residual activity of the factor remains even when the ligand is used in a molar excess. A curve may be established of the inhibiting effect of a ligand in accordance with the present invention with respect to a certain targeted factor against the concentration of the said ligand and the concentration may be determined at which a minimal residual factor activity still exists which is at least 1%, preferably at least 2%. The residual factor activity at five times this concentration should not be substantially different from the residual activity at the minimal point. It is especially a further aspect of the present invention to provide high affinity monoclonal antibodies, both human and humanized, as well as fragments, derivatives, and homologs of any of these, having the capacity to only partially inactivate a factor or factors in hemostasis, in particular in the coagulation cascade and more in particular factor VIII or a complex thereof, even in molar excess of the ligand, thereby preventing the risk of overdosage and the resulting bleeding complications. It is still another aspect of the present invention to provide a novel cell line producing the respective human monoclonal antibody.

The present invention also includes polynucleotide sequences which encode for the antibodies or fragments thereof mentioned above. It will be appreciated that a multitude of nucleotide sequences exist which fall under the scope of the present invention as a result of the redundancy in the genetic code. The present invention also includes complementary sequences which correspond to the monoclonal antibodies, or fragments thereof, mentioned above. In particular, the present invention includes probes constructed from the monoclonal antibodies, or fragments thereof, mentioned above or from the polynucleotides or from the complementary sequences mentioned above.

The present invention further provides a method of attenuation of coagulation in humans, comprising administering a ligand, being other than a polyclonal antibody,

such as a monoclonal antibody, either human or humanized, fragment, derivative or homolog thereof, capable of only partially inactivating a factor or factors in hemostasis, in particular in the coagulation cascade and more in particular factor VIII or a complex thereof to a patient in need of such attenuation even when the said ligand is in a molar excess. It further provides a method of treatment or prevention of a thrombotic pathologic condition in mammals, namely in humans, comprising administering a therapeutically effective amount of a ligand, other than a polyclonal antibody, for instance a monoclonal antibody, either human or humanized, or a fragment, derivative or homolog thereof, capable of only partially inactivating, even when the said ligand is in a molar excess, a factor or factors involved in hemostasis, in particular in the coagulation cascade, and more particularly factor VIII or a complex including factor VIII, to a mammal in need of such treatment or prevention. In a preferred embodiment, the thrombotic pathologic condition may be selected for instance from intravascular coagulation, arterial thrombosis, arterial restenosis, venous thrombosis and arteriosclerosis.

Another embodiment of the present invention is directed to a pharmaceutical composition comprising a ligand, other than a polyclonal antibody, having the capacity of binding to a site on a factor or factors involved in hemostasis, in particular in the coagulation cascade, and more particularly factor VIII or a complex including factor VIII, for only partially inactivating the said factor or factor complex even when the ligand is in molar excess, in admixture with a pharmaceutically acceptable carrier. The said ligand preferably is a high affinity anti-factor VIII or anti-factor VIII - von Willebrand factor complex monoclonal antibody, either human or humanized, or hybridized, or a fragment, derivative or homolog thereof. The pharmaceutical composition of the present invention may further optionally comprise a therapeutically effective amount of a thrombolytic agent.

Another embodiment of the present invention is directed to methods for the selection of specific monoclonal antibodies. The conventional technique of immunizing an animal such as a mouse with a protein such as factor VIII elicits an immunological response which may involve several epitopes on the factor VIII molecule. The present invention provides more selective methods of obtaining specific monoclonal antibodies against an epitope of a wild-type protein. First, a donor, e.g. a mammal such as a human, is provided (i.e. selected) which has an at least partially functional modified version of a wild-type protein. The said modification, which lies in a domain of the protein, may be due to any cause, e.g. race or variety, to genetic defects at birth, to an illness or by human interference, e.g. immunotolerance against the functionally modified version. The mammal donor is then administered the wild-type protein in order to elicit an immune response; at this stage, it is important that a sufficient quantity

of the wild-type protein (e.g. factor VIII) be administered until an immune response is generated. Then, in a final step of the method, selection of B-cells from the mammal donor will result in a much greater chance of obtaining monoclonal antibodies against an epitope in the region of the modification, for instance by selecting B-lymphocytes from the donor which produce antibodies only partially inactivating the wild type protein.

The anticoagulant potential of inhibiting factor VIII has to date not been explored, perhaps because of the well known bleeding complications that occur in hemophilia A patients that lack factor VIII activity completely (severe hemophilia) or to a large extent (moderate hemophilia). Hemophilia A, however, not only demonstrates the importance of factor VIII as limiting co-factor of coagulation, but also the existing link between coagulation and the development of atherosclerosis. Atherosclerosis and its thrombotic complications were indeed found to be significantly rarer among patients with hemophilia A. Antagonizing factor VIII activity at a level that allows sufficient hemostasis to prevent bleeding but protects from pathologic intravascular thrombus formation therefore holds substantial promise for safe anticoagulation in prothrombotic diseases such as deep vein thrombosis (DVT), pulmonary embolism (PE), postoperatively, in pregnancy, in coronary artery disease (CAD), cerebrovascular disease (CVD), peripheral artery disease (PAD) and in vascular interventions.

The present invention is based on on the surprising determination of new ligands, namely new human and humanized monoclonal antibodies and fragments, derivatives and homologs thereof. These may exhibit an unforeseen "plateau effect", i.e. the achievement of only a partial inactivation of a factor involved in hemostasis, in particular in the coagulation cascade, either individually or in combination, whatever the excess of ligand. The ligands may bind to a factor or a complex of factors resulting in only partially impairing the function of a physiologically functional site of the said factor or factor complex. This "plateau effect" makes the ligands particularly suitable for treating coagulation disorders and resulting thrombotic pathologic conditions while minimizing the risk of bleeding, by comparison, notably, to antibodies with self-limiting neutralizing activity mentioned in WO97/26010. There is therefore a sharp contrast between the "the self-limiting neutralizing activity of antibodies to coagulation factors disclosed in WO97/26010 and the clinically meaningful plateau inhibition that the present invention covers, where anti- factor VIII antibodies such as KR1X-1 inhibit factor VIII activity by no more than 85%.

Particularly useful is a property of ligands in accordance with the present invention to allow some physiological function of the affected site even when the ligand is in molar excess. The ligands may be anti-factor VIII antibodies or antibodies against a factor VIII complex, in particular human or human hybrid monoclonal antibodies

which bind to factor VIII or a factor VIII complex and at least partially inhibit the activity of factor VIII. Data indicate that type II inhibitors react with different antigenic determinants than type I antibodies and that these determinants are partially blocked in the factor VIII/von Willebrand factor complex.

5       The present invention will now be described in more details with reference to the following drawings.

#### BRIEF DESCRIPTION OF THE DRAWINGS

10       Fig. 1 presents the results of producing human monoclonal antibodies derived from a hemophilia A patient, expressed in the form of IgG antibody binding to factor VIII in ELISA.

Fig. 2 shows inhibition of factor VIII activity by the monoclonal antibody BO2C11.

15       Fig. 3 shows inhibition of factor VIII activity by the monoclonal antibody produced from the cell line KRIX 1.

Fig. 4 shows inhibition of the binding of activated factor VIII on phosphatidyl-L-serine by the monoclonal antibody produced from the cell line KRIX 1.

Fig. 5 shows the influence of certain polyclonal antibodies on the dissociation of activated factor VIII from von Willebrand factor.

20       Figs. 6 and 8 show amino acid sequences (the lower lines) and nucleotide sequences (upper lines) for the variable regions  $V_H$  of the heavy chains of BO2C11 and the KRIX 1 monoclonal antibodies, respectively. Also shown are the three complementarity determining regions (CDR) of each chain which are each an individual polypeptide ligand in accordance with an individual embodiment of the present invention.

25       Figs. 7 and 9 show amino acid sequences (the lower lines) and nucleotide sequences (upper lines) for the variable regions  $V_L$  of the light chains of BO2C11 and the KRIX 1 monoclonal antibodies, respectively. Also shown are the three CDR's of each chain which are each an individual polypeptide ligand in accordance with an individual embodiment of the present invention.

30       Fig. 10 provides a graph showing inhibition of factor VIII activity by the antibody SAF8C-Ig mentioned in WO97/26010.

Fig. 11 illustrates the kinetics of factor VIII inhibition by KRIX-1.

35       Fig. 12 illustrates the inhibition of venous thrombosis in a hamsters model by KRIX-1.

## DEFINITIONS

The term "antibody" refers to intact molecules as well as fragments thereof, such as Fab, Fab', F(ab')<sub>2</sub> or Fv, which are capable of binding to the epitope determinant of the relevant factor or domain of the factor.

5 "Humanized antibody" as used herein, refers to antibody molecules in which amino acids have been replaced in the non-antigen binding regions in order to more closely resemble a human antibody.

A "Reshaped human antibody" or a "Human hybrid antibody" as used herein, refers to a human antibody in which amino acids in the antigen binding regions have  
10 been replaced with sequences in accordance with the present invention, e.g. CDR's, or other parts of variable regions which have been derived from the repertoire of human antibodies.

The term "homology" or "homologous" as used herein with reference to ligands in accordance with the present invention refers to a molecule which will compete with  
15 or inhibit binding of one of the ligands in accordance with the present invention to the target site. The binding should be specific, i.e. the binding of the alternative molecule should be as specific to the site as the ligand in accordance with the present invention. Where the ligands in accordance with the present invention include amino acid sequences, homology may include having at least 80%, more preferably 90% and most  
20 preferably 95% amino acid sequence identity with the relevant ligand.

## DETAILED DESCRIPTION OF THE INVENTION

The present invention will be described with reference to certain embodiments and to certain figures but the present invention is not limited thereto but only by the  
25 claims. In particular, the present invention will mainly be described with reference to ligands to factor VIII but the present invention is not limited thereto. The present invention relates to a general concept of obtaining a therapeutically useful "plateau inhibition" by only partially inactivating a factor in hemostasis by the selection of certain monoclonal antibodies, as well as producing such human or humanized  
30 monoclonal antibodies, or fragments, derivatives or homologs thereof, and using these for anti-thrombotic therapy and in anti-thrombotic therapeutic compositions. These ligands and compositions may have the advantageous property that the inactivation of the factor is only partial even when the ligand is in a molar excess. This means that even though the ligand is used in an amount which might be expected to inactivate  
35 completely the targeted factor, the inactivation is still incomplete.

The present invention provides a particular cell line producing human monoclonal antibodies which are reactive with human factor VIII and more specifically have the capacity of inactivating the co-factor activity of human factor VIII by

interfering with proteolytic cleavage site or von Willebrand factor or tenase complex reaction or by inducing a three-dimensional conformational change in factor VIII, in particular by targeting a domain of factor VIII and by recognizing epitopes located in the said domain. One preferred domain is the C1 domain of factor VIII but the present invention is not limited thereto. A site on the C2 domain of factor VIII may also be partially inhibited. The present invention also includes ligands other than polyclonal antibodies, in particular monoclonal antibodies, which reduce the release rate of factor VIII from von Willebrand factor. These monoclonal antibodies specifically target factor VIII when bound to von Willebrand factor and hence target an epitope associated with the complex of factor VIII and von Willebrand factor. The present invention also provides fragments of any of the above monoclonal antibodies such as Fab, Fab', F(ab')<sub>2</sub>, scFv, CDR's, single variable domains as well as derivatives, homologs and combinations of these. More particularly, these monoclonal antibodies and fragments may target a domain of factor VIII, in particular the C1 domain of factor VIII. They may also partially inhibit a site on the C2 domain of factor VIII. They may also target an epitope associated with the complex of von Willebrand factor and factor VIII. An aspect of the present invention is therefore to provide ligands other than polyclonal antibodies which bind to a first site (e.g. in the C1 domain of factor VIII) remote from a functional second site (e.g. the site in the C2 domain of factor VIII which is responsible for binding phospholipids) in such a way that the function of the second site is only partially impaired even when the ligand is in a molar and therapeutic excess.

The cell line named KRIX 1 producing monoclonal antibodies according to the present invention was deposited with the BCCM/LMBP (Belgian Co-ordinated Collections of Microorganisms/Plasmid Collection Laboratorium voor Moleculaire Biologie, University of Ghent K.L. Ledeganckstraat 35, B-9000 Ghent, BE under accession number LMBP 5089CB on July 1, 1999.

The present invention further provides cell lines producing human monoclonal antibodies having a reactivity substantially similar to that of the human monoclonal antibodies obtained from the above-mentioned deposited cell line, as well as the human monoclonal antibodies obtainable from these further cell lines.

The present invention further provides reshaped human monoclonal antibodies or human hybrid monoclonal antibodies against factor VIII, which bind to and only partially inactivate factor VIII or a complex including factor VIII and von Willebrand factor which comprise only elements derived from the repertoire of human antibodies. By human hybrid monoclonal antibodies it is meant a hybrid antibody constructed from a human antibody and from variable regions in accordance with the present invention. Conventionally in the art, until now it has only been possible to obtain antibodies

against factor VIII derived from animals, e.g. mice, or to construct chimeric antibodies from human antibodies with the variable portions derived from mice antibodies.

The present invention also provides ligands, other than polyclonal antibodies, having the capacity of only partially inactivating a factor (or a complex including a factor) involved in hemostasis, in particular in the coagulation cascade of blood, preferably factor VIII or a complex including factor VIII, by binding to a site of the said factor or complex, the said only partial inactivation also taking place when the ligand of the invention is in a molar excess with respect to the said factor. The site to which the ligand binds may or may not be directly or substantially involved in a physiological interaction of the said factor or complex. For instance, the ligand may bind to a site which is at a predetermined distance away from a physiologically functional site of the said factor. By partial "plateau" inactivation, herein we mean an at most 98% inactivation, preferably an at most 95% inactivation, as determined by a suitable test method such as for instance the chromogenic assay available from Coatest® (Kabi Vitrum, Brussels, Belgium) or from Chromogenix AB, Mölndal (Sweden). The level of activation required may depend upon the physiological function of the factor involved in hemostasis. On the other hand, in order to provide therapeutic usefulness, inactivation of the blood factor should be at least about 65%, preferably at least about 70%, as determined by the same test method as above. It will be appreciated that the ligands in accordance with the present invention operate in a different way from the mechanism conventionally ascribed to type II antibodies against factor VIII. One conventional mechanism is that of competition with another factor, e.g. von Willebrand factor. The kinetics of a competition mechanism mean that if the one species is at a high concentration compared with the other (e.g. in a molar excess), the inhibition is effectively complete. In contrast, the ligands of the present invention reach a plateau in the inactivation of the relevant factor, which is substantially independent of the excess of the ligand. The other conventional mechanism ascribed to type II antibodies is that of low affinity: also in this case, an excess will drive the reaction to complete inhibition.

When the targeted blood factor is factor VIII, the ligands of the invention may be human monoclonal antibodies obtainable from the deposited cell line KRIX 1, preferably being of class IgG, and which have the capacity of only partially inactivating the co-factor activity of factor VIII. More specifically the invention relates, in a preferred embodiment, to human monoclonal antibodies from such origin which are able to recognize epitopes located in the C1 domain of factor VIII. Although the inventors do not wish to be bound to a single explanation or theory, it is believed that the binding of such human monoclonal antibodies results in partial impairment of the binding of activated factor VIII to phospholipids, a necessary step for cofactor activity expression.

The present invention further provides monoclonal antibodies having substantially the same characteristics as above disclosed and being produced by on purpose immunization in animals, preferably in mouse, for instance by injecting human factor VIII in mice and then fusing the spleen lymphocytes with a mouse myeloma cell line, followed by identifying and cloning the cell cultures producing anti-factor VIII antibodies. The monoclonal antibodies produced in animals are then humanized, for instance by associating the binding complementarity determining region ("CDR") from the non-human monoclonal antibody with human framework regions - in particular the constant C region of human gene - such as disclosed by Jones et al. in *Nature* (1986) 321:522 or Riechmann in *Nature* (1988) 332:323.

The present invention also provides fragments and derivatives, in particular complementarity determining regions ("CDR's") of the above monoclonal anti-factor VIII antibodies as well as homologs thereof. For instance, the invention provides antigen-binding fragments Fab, Fab' and F(ab')<sub>2</sub> generated by proteolytic digestion of the said monoclonal antibodies using methods well known in the art, such as described by Stanworth et al., *Handbook of Experimental Immunology* (1978), vol.1 chapter 8 (Blackwell Scientific Publications). Such fragments, which contain the antibody binding site, have lost a number of properties of the parent antibody, such as complement activation or capacity to bind to Fc gamma receptors. The present invention also includes single chain fragment variables (scFv), single variable domain fragments of the antibodies and combination of these fragments and of the above mentioned fragments.

The invention also provides soluble or membrane anchored single-chain variable parts of the above monoclonal antibodies and a method for their obtention as follows. The DNA sequences of the variable parts of human heavy and light chains are amplified in separated reactions and cloned. A fifteen amino-acid linker sequence, for instance (Gly4 Ser)<sub>3</sub>, is inserted between VH and VL by a two-steps polymerase chain reaction (PCR), for instance according to Dieffenbach and Dveksler, "PCR Primer, a laboratory manual" (1995), Cold Spring Harbour Press, Plainview, NY, USA. The resulting fragment is then inserted into a suitable vector for expression of single chain fragment variable (scFv) as soluble or phage-displayed polypeptide. This can be achieved by methods well known to those skilled in the art, such as described by Gilliland et al., *Tissue Antigens* (1996) 47:1-20. The present invention also includes a ligand comprising peptides representative of hypervariable regions of a monoclonal antibody which can be obtained by synthesis using an applied biosystem synthesizer, for instance a polypeptide synthesizer such as model 9050 available from Milligen (USA) or a model from a related technology, which alone or in combination with other or similar hypervariable regions will exert properties similar to that of the parent antibody.



The invention further provides a pharmaceutical composition for the prevention or treatment of disorders of hemostasis, in particular of the coagulation cascade and resulting thrombotic pathologic conditions in humans, comprising, as an active ingredient, a ligand other than a polyclonal antibody, preferably a human monoclonal antibody such as disclosed hereinabove, in admixture with a pharmaceutically acceptable carrier. More preferably the said monoclonal antibody is a human monoclonal antibody, or a fragment, derivative or homolog thereof, obtainable from the cell line KRIX 1 deposited with the Belgian Co-ordinated Collections of Micro-organisms under accession number LMBP 5089CB. The degree of homology with the said monoclonal antibody is preferably at least 80%, more preferably 90% and most preferably 95%, and the homology is preferably particularly in respect to the complementarity determining regions of the antibody. A ligand in accordance with the present invention may also include a synthetic polypeptide of equivalent potency. The pharmaceutical composition of the present invention should comprise a therapeutically effective amount of the said above ingredient, such as indicated hereinafter in respect to the method of treatment or prevention.

The pharmaceutical composition of the present invention may further comprise, namely in view of a so-called adjunctive anti-thrombotic treatment, a therapeutically effective amount of a thrombolytic agent. Such thrombolytic agents, as well as their usual dosage depending on the class to which they belong, are well known to those skilled in the art. Among numerous examples of thrombolytic agents which may be included in the pharmaceutical compositions of the invention, the following non-limiting list may be particularly cited: t-PA, streptokinase, reptilase, TNK-t-PA or staphylokinase.

Suitable pharmaceutical carriers for use in the pharmaceutical compositions of the invention are described for instance in Remington's Pharmaceutical Sciences 16<sup>th</sup> ed. (1980) and their formulation is well known to those skilled in the art. They include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents (for example phenol, sorbic acid, chlorobutanol), isotonic agents (such as sugars or sodium chloride) and the like. Additional ingredients may be included in order to control the duration of action of the monoclonal antibody active ingredient in the composition. Control release compositions may thus be achieved by selecting appropriate polymer carriers such as for example polyesters, polyamino acids, polyvinyl pyrrolidone, ethylene-vinyl acetate copolymers, methylcellulose, carboxymethylcellulose, protamine sulfate and the like. The rate of drug release and duration of action may also be controlled by incorporating the monoclonal antibody active ingredient into particles, e.g. microcapsules, of a polymeric substance such as hydrogels, polylactic acid, hydroxymethylcellulose, polymethyl methacrylate and the other above-described

polymers. Such methods include colloid drug delivery systems like liposomes, microspheres, microemulsions, nanoparticles, nanocapsules and so on. Depending on the route of administration, the pharmaceutical composition comprising the active ingredient may require protective coatings. The pharmaceutical form suitable for injectionable use include sterile aqueous solutions or dispersions and sterile powders for the extemporaneous preparation thereof. Typical carriers therefore include biocompatible aqueous buffers, ethanol, glycerol, propylene glycol, polyethylene glycol and mixtures thereof.

The present invention also provides the use of a ligand, other than a polyclonal antibody (as disclosed above) as a medicament. More preferably the medicament used in the present invention is a means for preventing and/or treating disorders of hemostasis, in particular, coagulation disorders and other thrombotic pathologic conditions in mammals, preferably in humans. The said ligand may be provided to a patient by any means well known in the art, i.e. orally, intranasally, subcutaneously, intramuscularly, intradermally, intravenously, intraarterially, parenterally or by catheterization. According to the present invention, the ligand may also be used as a medicament in conjunction or association with another medicament, for instance a thrombolytic agent such as disclosed hereinabove under the heading of pharmaceutical compositions.

The present invention therefore provides a method of treatment and/or prevention of hemostasis, coagulation disorder or thrombotic pathologic condition as well as a method of attenuation of coagulation in a mammal, preferably a human, comprising administering to a mammal in need of such treatment or prevention or attenuation of coagulation a therapeutically effective amount of a ligand other than a polyclonal antibody such as disclosed hereinabove. Preferably the said ligand is a human or humanized monoclonal antibody obtainable from cell line KR1X 1 deposited with the Belgian Co-ordinated Collections of Micro-organisms under accession number LMBP 5089CB or an antigen-binding fragment Fab, Fab' or F(ab')<sub>2</sub>, a complementarity determining region (CDR), a soluble or membrane-anchored single-chain variable part (scFv), a single variable domain or a derivative or combination of any of these elements.

A therapeutically effective amount as used herein means from about 1 microgram to about 10 milligrams per kilogram of body weight, more preferably from about 10 micrograms to about 1 milligram per kilogram of body weight of the mammal to be treated. It will be appreciated that, in view of the long half-life time of most IgG human antibodies, the ligands of the present invention which are monoclonal antibodies of the said class will enjoy a periodicity of treatment which participates in the comfort of the patient.

As preferred embodiments of the said thrombotic pathologic condition to be prevented or treated, may be cited intravascular coagulation, arterial thrombosis (which may be responsible for acute myocardial infarction and stroke), arterial restenosis, venous thrombosis (which commonly occurs in peripheral veins as a consequence of accidental or surgical trauma or immobilization) or arteriosclerosis. In a most preferred method of treatment, the patient is provided with a bolus (intravenously injected) at a dosage determined by the ordinary skilled physician depending on criteria which establish the particular patient's clinical condition.

The method of treatment and/or prevention according to the invention may include further treatment or prevention of the same thrombotic pathologic condition by administering, preferably by sequentially administering, to the patient a therapeutically effective amount of a thrombolytic agent such as disclosed hereinabove under the heading of pharmaceutical compositions. Sequentially, as used herein, means that the ligand of the present invention and the known thrombolytic agent are administered to the patient sequentially but not simultaneously.

The present invention further provides a method of obtaining monoclonal antibodies from a non-human mammal, comprising the steps of:

- a) selecting a non-human mammal having a modified and at least partially functional physiologically active protein, the modification being with respect to a wild type protein and lying in a domain of the protein;
- b) administering the wild type protein to the non-human mammal in order to elicit an immune response, and
- c) selecting B-lymphocytes from the non-human mammal which produce antibodies which only partially inactivate the wild type protein.

According to this method, the standard practice is to sacrifice the non-human mammal and to remove its spleen in order to perform step (c).

The present invention additionally provides a method of obtaining monoclonal antibodies from the blood of a human being having a modified and at least partially functional physiologically active protein, the modification being with respect to a wild type protein and lying in a domain of the protein, and to whom the wild type protein was administered, the said method comprising the step of selecting, from the blood of said human being, B-lymphocytes which produce antibodies which only partially inactivate the wild type protein.

The present invention, as embodied in the various above disclosed aspects, has a number of advantages. The major advantage of the therapeutic use of the human monoclonal antibodies of the invention is that the treatment is highly specific for the immune response under consideration. In hypercoagulation states, the specificity of the

human monoclonal antibodies of the invention ensures that interaction within the coagulation cascade pathway is limited to the factor recognized by the antibody.

More specifically, the use of the anti-factor VIII antibodies having the above-mentioned preferred characteristics brings a unique combination of the advantages related to the targeting of factor VIII, those related to the characteristics of factor VIII inhibition and those related to the use of antibodies:

- targeting factor VIII means that neutralizing a co-factor activity such as that of factor VIII carries no risk of completely inhibiting the enzymatic activity it enhances, thereby representing an advantage over methods targeting directly enzymes such as factor IX.

- the embodiments of the inhibitors described above have in common that they efficiently but only partially inhibit the co-factor activity of factor VIII, setting a therapeutically useful plateau, even when the monoclonal antibody of the invention is used in more than 100-fold excess. Monoclonal antibodies in accordance with the present invention achieve a plateau effect in inactivation of factor VIII, allowing bolus application, yielding safe antithrombotic protection over several weeks without the need of monitoring or the risk of overdosage.

- human IgG antibodies exhibit a prolonged half-life time of three weeks (except for IgG3 which is one week), thus providing very stable plasma levels of the anti-thrombotic agent and allowing for a drastic reduction in the frequency of administration. Further, the use of human antibodies or derivatives carries a minimal risk of inducing immune responses.

The present invention is further described by the following examples which are provided for illustrative purposes only.

#### EXAMPLE 1 - production of monoclonal antibodies derived from hemophilia A patients.

Human monoclonal antibodies of the desired specificity and characteristics are produced by transformation of B lymphocytes obtained from the peripheral blood of patients suffering from hemophilia A or acquired hemophilia. The method of selecting patients is an embodiment of the present invention. In order to elicit a more specific immunological response, patients are sought who have an impaired function of a physiologically active protein, e.g. factor VIII. By "impaired" is meant that some residual function is available but that this is less than is known for the wild-type of the same protein. A comparison between the self-protein and the wild-type protein should exhibit a difference in the two proteins, preferably in a region or domain which is of interest. The difference may be a deletion or a substitution of one or more amino acids with others. The patients are then administered enough of the wild-type protein to elicit

an immunological response. Then, B-lymphocytes are extracted from the patients and selected based on the production of antibodies which have desirable properties. Although reference is made to "patients" above, the method in accordance with this embodiment may be applied generally to mammals. The above procedure results in a greater chance of obtaining antibodies which target the domain containing the defect.

B cells are transformed by infection with the Epstein-Barr virus and activation of surface antigens using techniques well known by those skilled in the art. Cell supernatants containing appropriate antibodies are identified by a specific assay procedure such as described in more details hereinbelow.

Thus, antibodies towards factor VIII are identified by reacting the supernatant with polystyrene microtitration plates coated with factor VIII or with factor VIII/von Willebrand factor complexes. The binding of specific antibodies is detected by addition of a non human IgG reagent coupled to an enzyme. Addition of an enzyme substrate which is converted to a colored compound in the presence of the enzyme allows the detection of specific antibodies. Such methods referred to as enzyme-linked immunoassays (ELISA) are well known to those skilled in the art and described in details e.g. in *Current Protocols in Immunology*, chapter 2, John Wiley & Sons (1994), the content of which is incorporated herein by reference.

More specifically in the present case, the binding of anti-factor VIII IgG antibodies was detected by addition of a horseradish peroxidase labeled mouse monoclonal antibody specific for human Fc gamma. The IgG subclass of the anti-factor VIII antibody was detected in ELISA, as presented in Fig. 1. The inhibition of factor VIII functional activity was tested in a functional coagulation assay as follows. Equal volume of cell culture supernatant and of a pool of normal plasma were incubated for two hours at 37°C and the residual factor VIII activity measured thereafter. Those antibodies which significantly inhibit factor VIII activity are shown with an asterisk in Fig. 1.

B cells (such as BO 2C11) producing anti-factor VIII antibodies are then expanded and cloned by limiting dilution as described for instance in *Current Protocols in Immunology* (see supra). Anti-factor VIII antibodies having the capacity to inhibit the procoagulant activity of factor VIII as described above are identified using a chromogenic assay kit such as a factor VIII chromogenic assay from Dade, Dürchingen, Germany or Coatest® commercially available from Kabi Vitrum (Brussels, Belgium) or Chromogenix AB (Mölnä, Sweden).

Equal volumes of monoclonal antibodies BO 2C11 and a pool of normal blood plasma were incubated for 2 hours at 37°C. BO 2C11 concentrations before mixing are shown on the X axis. The reduction of factor VIII activity was measured in a coagulation assay and was expressed as a percentage of the activity obtained in the

absence of antibody (see Fig. 2). The residual activity goes to zero asymptotically (complete inhibition).

Antibodies which inhibit factor VIII function with sufficient affinity but do not inhibit factor VIII pro-coagulant activity completely, even when used in large antibody excess, were selected in a further embodiment of the present invention. A representative example of such an antibody is provided in Fig. 3 where, equal volumes of KRIX 1 and of recombinant factor VIII or of normal plasma being incubated for two hours at 37°C and concentrations (expressed in microgr/ml) of KRIX 1 before mixing with plasma being as indicated, the residual factor VIII activity was measured using the above-mentioned chromogenic assay. Fig. 3 interestingly shows about 60 % factor VIII inhibition at a concentration of 0.1 microgr/ml and more interestingly an asymptotic factor VIII inhibition of about 80% in the whole range of concentrations from 0.5 to 300 microgr/ml.

The thus selected antibodies are then produced in bulk culture and purified by affinity chromatography using methods well known to those skilled in the art.

The details of a non-limiting preparation technique are as follows. Human recombinant factor VIII (specific activity: 4000 IU/mg) was obtained from Hyland (Glendale, Ca) as material for laboratory use only; plasma-derived (pd) factor VIII-von Willebrand factor complex, purified by ion exchange chromatography (specific activity  $\pm 160$  IU/mg protein; 15:1 von Willebrand factor to factor VIII w/w ratio), and purified factor VIII-depleted von Willebrand factor (von Willebrand factor to factor VIII w/w ratio 4800:1; lot 951016) were obtained from the Belgian Red Cross (Brussels, Belgium).

Peripheral blood samples were collected from donors suffering mild hemophilia and with inhibitors. The peripheral blood mononuclear cells (PBMC) were immortalized by EBV infection concomitantly to the activation of surface antigens. Four hundred and eighty cell lines were screened by ELISA for production of antibodies towards factor VIII. For example, one cell line, named KRIX 1, was successfully cloned by limiting dilution. Clonality was verified by RT-PCR amplification of mRNA coding for the V regions of the antibody heavy and light chains: a single sequence was obtained from 10 clones of PCR products. Purified antibodies were obtained by passage of KRIX 1 cell culture supernatant on Protein-A Sepharose. An ELISA performed with IgG subclass- and light chain-specific antibodies identified KRIX-1 as an IgG4k.

Human monoclonal antibodies were purified by adsorption on immobilized Protein A (high-TRAP<sup>R</sup> Protein A; Pharmacia, Uppsåla, Sweden). Fab fragments of human monoclonal antibody were prepared by papain digestion. One mg of a selected antibody was diluted to 500 microgr/ml in phosphate buffer (40 mmol/L KH<sub>2</sub>PO<sub>4</sub>, 60

mM Na<sub>2</sub>HPO<sub>4</sub>·2H<sub>2</sub>O, 0.15M NaCl) containing 50 mmol/L L-cystein (Sigma), 1 mmol/L EDTA (Merck) and 10 microgr papain (Sigma). The mixture was incubated for 3 h at 37°C with continuous agitation. The reaction was stopped by addition of iodoacetamide to a final concentration of 75 mmol/L for 30 min at RT. The digested antibody was dialysed against phosphate-buffered saline (140 mmol/L NaCl, 67 mmol/L KCl, 20 mmol/L Na<sub>2</sub>HPO<sub>4</sub>, 4.4 mmol/L KH<sub>2</sub>PO<sub>4</sub>, pH 7.4). The undigested IgG and Fc fragments were then eliminated by passage over protein A sepharose (Hi Trap Protein A; Pharmacia). The Fab fragment was further purified by gel filtration chromatography on a Superdex 200 (Pharmacia).

Conventional methods were used for the detection of anti-factor VIII IgG antibodies, the determination of IgG subclass, and the evaluation of inhibition of factor VIII binding to von Willebrand factor. For the analysis of the inhibition of the binding of rfactor VIII to a selected antibody by Fab and native antibody, Maxisorb polystyrene plates (Nunc) were coated for 2 h with 50 µl of the antibody diluted to 5 microgr/ml in glycine-buffered saline (20 mmol/L glycine, 34 mmol/L NaCl, pH 9.2). After washing, 50 µl of biotin-labeled rfactor VIII diluted to 1 microgr/ml in Tris-casein (10 mmol/L tris(hydroxymethyl)-aminoethane, pH 7.3, containing 150 mmol/L NaCl and 0.5% casein) were mixed for 1 h at 37°C with 50 µl of human IgG at various dilutions. A 50-µl aliquot of the mixture was added to the plates for 2 h at RT. After washing, the binding of biotinylated rfactor VIII was detected by sequential addition of avidin-peroxydase and OPD.

rfactor VIII (final concentration 0.2 microgr/mL) was incubated with human IgG antibody at different concentrations for 2 hours at 37°C and the residual factor VIII activity was assessed by a chromogenic assay (Coatest Factor VIII, Chromogenix AB, Mölndal, Sweden or Kabi Vitrum, Brussels, Belgium). Inhibition of plasma factor VIII activity was measured by the Bethesda method, in which a pool of normal plasma collected in buffered trisodium citrate was used as factor VIII source. Residual factor VIII activity was assessed by a chromogenic or by a one-stage clotting assay.

#### EXAMPLE 2 - Production of monoclonal antibodies by immunization in animals.

Alternatively, monoclonal antibodies having the same characteristics as disclosed in example 1 can be produced by on purpose immunization in animals. Thus, mice are injected with human factor VIII in Freund's adjuvant

Monoclonal anti-human factor VIII antibodies are then obtained by fusion of spleen lymphocytes with a mouse myeloma cell line. Cell culture supernatants producing anti-factor VIII antibodies are identified and cloned by limiting dilution, using methods described in *Current Protocols in Immunology* (see supra). Further

selection of inhibitors having the desired capacity to inhibit the pro-coagulant activity of factor VIII is carried out as described in example 1.

Monoclonal antibodies produced in mice are then humanized. Thus, sequences of the variable parts of mouse heavy and light chains are aligned with human immunoglobulin variable regions to identify human antibody with the greatest homology in framework regions. The DNA fragment encoding humanized variable regions are then synthesized by a PCR-based CDR (complementarity determining regions) grafting method as described for instance in Sato et al., *Cancer Research* (1993) 53:851-6. The final PCR product coding for the heavy chain variable part of the humanized antibody is digested and subcloned upstream of the human C gamma-1 gene in a first expression plasmid. The humanized light chain variable region of the final construction is inserted upstream of the C kappa gene in a second expression plasmid. The two constructions are then co-transfected into COS cells expression system.

### 15 EXAMPLE 3 - Characterization of anti-factor VIII antibodies.

Monoclonal antibodies of either human (example 1) or animal (example 2) origin are characterized using an assay system by which their capacity to inhibit the binding of factor VIII to phospholipids is evaluated. Thus, polystyrene microtitration plates are coated with phosphatidyl-L-serine. Soluble recombinant factor VIII at 2 microgr/ml final concentration is mixed for 30 minutes at 37°C with various concentrations of the antibody under evaluation. The mixture is then rapidly activated with thrombin and added to phosphatidyl-L-serine coated plates. The said plates were then incubated for two minutes at 21°C and the binding of factor VIII was detected by addition of the anti-factor VIII A1 domain mAbF14A2, for two minutes, followed by a two minute incubation with HRP-conjugated goat anti-mouse Fc gamma. Results of this experiment are shown in Fig. 4 for the monoclonal antibody produced from the cell line KRIX 1. On the figure, the average of activated factor VIII binding in the absence (closed symbols) or presence (open symbols) of the antibody, as well as the standard deviation of triplicates are indicated. Controls in the absence of factor VIII gave OD 490 lower than 0.05. Fig. 4 clearly shows that the monoclonal antibody produced from cell line KRIX 1 inhibits significantly the binding of factor VIII to phospholipids but brings about incomplete inhibition even when added in large excess.

To demonstrate that in absence of plasma, KRIX 1 did not recognize the mutated factor VIII light chain of the donor, DNA fragments encoding wild-type and mutated factor VIII light chains were synthesized. The corresponding proteins were expressed in reticulocyte lysates. The correct folding of native and mutated light chains was determined by immunoprecipitation with the human monoclonal antibody BO2C11, which recognizes a conformational epitope within the carboxy-terminal part



of the factor VIII light chain. Immunoprecipitation experiments indicated that BO2C11 bound wild-type and Arg2150His light chains, whereas KRIX 1 captured exclusively the wild-type light chain. Prolonged exposure of SDS-PAGE gels to the autoradiography film failed to detect any significant binding of KRIX 1 to the mutated light chain. Control experiments showed no binding to assay reagents other than factor VIII or factor VIII fragments, and preincubation with soluble rfactor VIII prevented the binding to methionine-labeled factor VIII fragments, confirming the binding specificity.

KRIX 1 did not recognize factor VIII in Western blotting indicating that the epitope recognized was conformational. Further epitope mapping was therefore performed with factor VIII fragments produced in reticulocyte lysates. Preliminary experiments had indicated that such an approach was efficient for the synthesis of factor VIII domains. The immunoprecipitation procedure using labeled factor VIII domains produced in reticulocyte lysate was validated by mapping the epitope recognized by the human monoclonal antibody BO2C11. A complete agreement was observed between the binding to factor VIII C2 deletion fragments produced in reticulocyte lysates and the binding to recombinant fragments produced in *E. coli* or COS cells. KRIX 1 bound to full-length light chain, to fragments corresponding to A3C1, C1C2, and the isolated C1 domain. In contrast, KRIX 1 did not bind to the C1 or C1C2 domains with the substitution Arg2150His, although in a control experiment, the Arg2150His C1C2 domain was bound by BO2C11 like its normal counterpart.

A search was made for other mutations in the light chain which could alter the binding of KRIX 1. As shown in Table I, KRIX 1 inhibited the activity of all mutated factor VIII molecules tested so far, except those carrying the mutation Arg2150His.

Table I. Inhibition of factor VIII activity in mild hemophilia A patients' plasma.

factor VIII mutation	factor VIII activity (IU/ml)	KRIX-1 Inhibition (%)
Arg1689Leu	0.04	>70
Arg1749His	0.39	>70
Gly1750Arg	0.38	>70
Ala1824Val	0.16	>70
Asp1825Gly	0.17	>70
His1961Tyr	0.24	>70
Arg1966Gln	0.08	>70
Met2010Ile	0.04	>70

Ser2011Asn	0.23	>70
Val2016Ala	0.05	>70
Asn2019ser	0.13	>70
Leu2052Phe	0.23	>70
Asp2074Gly	0.13	>70
Thr2086Ile	0.08	>70
Ile2098Ser	0.20	>70
Phe2101Leu	0.06	>70
Asn2129Ser	0.20	>70
Arg2150His	0.03	0
Pro2153Gln	0.02	65
Arg2159Leu	0.16	>70
Trp2229Cys	0.02	>70
Gln2246Arg	0.08	>70

In the use of KRIX 1 as a medicament to partially inhibit factor VIII mild mutations of factor VIII will not affect the effectiveness of the therapy.

KRIX 1 inhibited the binding of factor VIII to von Willebrand factor in a dose-dependent manner. The concentration of KRIX 1 required to achieve 50% inhibition (IC<sub>50</sub>) of factor VIII binding was 0.25 microgr/mL and more than 95% inhibition was obtained with 20 microgr/mL of KRIX 1. Fab fragments of KRIX 1 also inhibited factor VIII binding to von Willebrand factor. However, on a molar basis 15 times more Fab than native KRIX 1 was required to inhibit 50% of factor VIII binding to von Willebrand factor. Additional experiments were performed to exclude that the KRIX 1 Fab fragments still contained intact or partially digested antibody. SDS-PAGE analysis of the Fab fragments purified by protein A adsorption and gel filtration chromatography showed a single band. The presence of trace amounts of Fc gamma fragments remaining bound to Fab fragments was excluded in ELISA. Wells with insolubilised factor VIII were incubated with native or Fab KRIX 1. Binding of both Fab and native KRIX 1 could be detected by addition of peroxidase-labeled anti-kappa light chain IgG. By contrast, addition of peroxidase-labeled anti-Fc gamma IgG did not reveal any specific binding even when 100 microgr/ml of the Fab preparation were incubated into the

5 wells. By comparison, the addition 0.1 microgr/ml of the native antibody gave rise to a significant binding. In ELISA, a 15-fold higher concentration of Fab than of native antibody was required to inhibit 50% of the binding of biotinylated KRIX 1 onto insolubilised factor VIII, indicating that the Fab KRIX 1 fragment had a lower affinity for factor VIII than the native antibody. Accordingly, the requirement for higher concentrations of KRIX 1 Fab than of native antibody to inhibit factor VIII binding to von Willebrand factor should be attributed to the reduced affinity of KRIX 1 Fab fragments for factor VIII.

10 To determine whether KRIX 1 was representative of the polyclonal antibodies from the donor, a competitive assay was used. The binding of biotinylated KRIX 1 to insolubilised factor VIII was measured in presence of increasing concentrations of either KRIX 1, polyclonal IgG from the donor, or control polyclonal IgG. IgG from the donor dose-dependently inhibited KRIX 1 binding to factor VIII. The concentration of KRIX 1 and IgG from the donor inhibiting 50% of biotinylated KRIX 1 on factor VIII  
15 were of 0.3 microgr/ml and of 170 microgr/ml, respectively, whereas no inhibition was observed with the control IgG.

EXAMPLE 4 - Production of monoclonal antibodies derived from hemophilia A patients and which bind to the factor VIII - von Willebrand factor complex.

20 Alternatively, antibodies which reduce the release rate of factor VIII from von Willebrand factor are identified as follows. Polystyrene microtitration plates are coated with a specific antibody to von Willebrand factor. A solution of biotinylated recombinant factor VIII (0.5 microgr/mL) complexed to von Willebrand factor (5 microgr/mL) is mixed with various concentrations of IgG from a donor (Fig. 5 solid squares), e.g. the same patient as described above (from which KRIX 1 was derived),  
25 MoAb4H1D7, or of IgG from a non-hemophiliac subject (Fig. 5 solid triangles). IgG at the indicated concentrations was added to microtitration plates coated with a mouse antibody, MoAb4H1D7 against von Willebrand factor for an incubation of two hours at room temperature. After washing, factor VIII was activated by thrombin during two  
30 minutes at 37°C. Factor VIII bound to von Willebrand factor was detected by addition of avidine peroxidase. Controls included the detection of bound biotinylated factor VIII in the absence of thrombin digestion (OD450=460+47.7SD) and of biotinylated recombinant factor VIII after thrombin digestion in the absence of antibody (OD450=160+16.0SD).

35 Results of these experiments are shown in Fig. 5 for polyclonal antibodies. On this figure, the average values as well as the standard deviation of triplicates are indicated. Fig. 5 clearly shows that a significantly higher proportion of activated factor VIII remains bound to the plate in the presence of increasing concentrations of the

antibody, i.e. it demonstrates a reduction of the dissociation of activated factor VIII from von Willebrand factor in the presence of an inhibitor antibody recognizing factor VIII bound to von Willebrand factor. Monoclonal antibodies have been obtained from these polyclonal antibodies in accordance with the methods described in this invention, thus indicating that the present invention may be extended to monoclonal antibodies and fragments and derivatives thereof which bind to Factor VIII/von Willebrand factor complexes.

#### EXAMPLE 5: Sequencing of antibody variable domains

Sequencing of antibody variable domains was carried out as follows. The isolation of RNA from EBV-immortalized human B-cell lines was performed using TRIzol Reagent according to the manufacturer's instructions (Life Technologies). CDNA was synthesized with the SuperScript preamplification system for first-strand cDNA synthesis. The cDNA encoding the heavy chain variable region genes (V<sub>H</sub>) was amplified by polymerase chain reaction (PCR) using primers specific for the leader sequence of the V<sub>H</sub> families and for the first exon of the C gamma region, as described (Bakkus et al, *Blood*, 80:2326, 1992) Annealing was performed at 60°C for 40 PCR cycles. PCR products of the appropriate size (460 bp) were isolated from 1.5% agarose gel and cloned using the TA Cloning Kit (Invitrogen BV, Leek, The Netherlands). A PCR screening using couples of primers corresponding to the V<sub>H</sub> gene family of interest was performed on cultures of randomly selected colonies. Plasmid DNA from positive colonies were isolated using Wizard Plus Minipreps (Promega, Menlo Park, CA) and sequenced in both directions with Sequenase (US Biochemical, Cleveland, OH), according to the manufacturer's instructions. Analysis of the variable gene sequences was made using the V BASE Sequence Directory (Tomlinson et al, MRC Centre for Protein Engineering, Cambridge, UK).

The complete sequences of the V<sub>H</sub> and V<sub>L</sub> of the antibody BO 2C11 described in example 1 were submitted to the EMBL Nucleotide Sequence Database under the accession numbers AJ224083 and AJ224084, respectively.

The amino acid sequences shown in figures 6 and 7 define the V<sub>H</sub> and V<sub>L</sub> regions of the antibody BO2C11 including the three CDR's for each of the heavy and light chains. Also given are the polynucleotide sequences which encode for these regions. SEQ.NOs. 2 and 3 are the amino acid sequences of the heavy and light chains of BO2C11, respectively while SEQ. NOs. 5 and 6 provide the polynucleotide sequences coding for these variable regions.

The amino acid sequences shown in figures 8 and 9 define the V<sub>H</sub> and V<sub>L</sub> regions of the antibody KRIX-1 including the three CDR's 1-3 for each of the short and long chains. Also given are the polynucleotide sequences which encode for these

regions. SEQ. NOs. 8 and 1 are the amino acid sequences of the heavy and light chains of KRIX-1, respectively while SEQ. NOs. 7 and 4 provide the polynucleotide sequences coding for these variable regions.

5 EXAMPLE 6 (COMPARATIVE) - Inhibition of factor VIII activity by the antibody SAF8C-Ig.

Levels of factor VIII are measured in a functional assay following an incubation period of two hours at 37°C with various concentrations of the antibody SAF8C-Ig, using the chromogenic assay described in example 1. As shown in figure 10, the residual factor VIII activity is reduced in a dose dependent manner. Already at 100 microgr/ml of SAF8C-Ig, residual factor VIII activity is less than 1% of the normal activity. Such low factor VIII levels expose the patient to a high risk of spontaneous bleedings, as is well known for instance from Levine, *Ann.NY Acad.Sci.*(1975) 240:201 and Gilbert, *Mount Sinai J.Med.*(1977) 44: 339.

15

EXAMPLE 7 - Inhibition of venous thrombosis in hamsters by KRIX 1.

Thrombosis was experimentally induced in the femoral vein of anesthetized hamsters, by injecting the dye rose-bengal in the jugular vein and by exposure of the femoral vein to the green light of a Xenon lamp for 4 minutes (*Kawazaki et al. Thromb Haemost* (1999) 81: 306-11). As a consequence of illumination of the vessel, the dye decomposes and generates radicals injuring the vessel endothelial cells. Thus, subendothelial structures are exposed to the blood circulation and thrombus formation is initiated. The amount of thrombus formed is measured via transillumination of the injured vessel (*Kawazaki et al. Thromb Haemost* (1999) 81: 306-11) and is quantified via the amount of white light being transilluminated through the vessel. As represented in figure 11, when this experiment is performed in control animals, the average thrombus size measured in 13 hamsters is  $220,000 \pm 32,575$  (mean  $\pm$  SEM) Arbitrary light Units (A.U.), whereas treatment of a group of 12 hamsters with KRIX-1 (400-800 microgr/kg, given as a bolus immediately prior to induction of thrombosis) reduced the mean thrombus size to  $122,000 \pm 27,100$  A.U. ( $p = 0.0188$ , Mann-Whitney test).

30 Additionally, the kinetics of factor VIII inhibition by KRIX-1 was analyzed *ex-vivo* as follows : hamsters were injected intravenously with KRIX-1 (1600 microgr/kg). Levels of factor VIII:c were measured in a chromogenic assay (*Coatest Factor VIII<sup>R</sup>* (Chromogenix AB, Mölndal, Sweden), and *Factor VIII Chromogenic Assay* (Dade, Düdingen, Switzerland)) using plasma collected before and at different periods of time after injection. Figure 12 shows that in these hamsters, factor VIII activity is reduced from 1.6 IU/ml to 0.3 IU/ml already 30 minutes after antibody injection, thus confirming that KRIX-1 only partially inhibits factor VIII.

CLAIMS

1. A cell line named KRIX 1 deposited with the Belgian Coordinated Collections of Micro-organisms, under accession number LMBP 5089CB.
2. A cell line producing human monoclonal antibodies having a reactivity substantially identical to that of the human monoclonal antibodies obtained from the cell line of claim 1.
3. A ligand, being other than a polyclonal antibody, having the capacity of only partially inactivating a factor or a complex of two or more factors involved in hemostasis when the said ligand is in a physiological excess by binding to a site of the said factor or the said complex.
4. A ligand according to claim 3, wherein the binding site of the ligand is not directly or substantially involved in a physiological interaction of the said factor or said complex.
5. A ligand according to claim 3, wherein the factor or the said complex is involved in the coagulation cascade of blood.
6. A ligand according to claim 5, wherein the factor is factor VIII or factor VIII is involved in the complex.
7. A ligand according to claim 6, having the capacity of inactivating the co -factor activity of factor VIII by interfering with a proteolytic cleavage site or the von Willebrand factor or the tenase complex reaction or by inducing a three-dimensional conformational change in factor VIII or by targeting a domain of factor VIII, in particular the C1 domain of factor VIII, or by targeting factor VIII in the factor VIII-von Willebrand factor complex.
8. A ligand according to claim 3, being a human monoclonal antibody obtainable from the cell lines of claim 1 or claim 2.
9. A ligand according to claim 8, being of class IgG.
10. A ligand according to any of the claims 3 to 9, being able to recognize epitopes located in the C1 domain of factor VIII.

11. A ligand according to any of claims 3 to 7, having substantially the same characteristics as a ligand according to claim 8 and being a monoclonal antibody produced by on purpose immunization in animals.
- 5 12. A ligand according to any of claims 3 to 11, having the capacity of binding to a site which is at a predetermined distance of a physiologically functional site of the said factor.
- 10 13. A humanized monoclonal antibody obtainable from the monoclonal antibody of claim 11.
14. An antigen-binding fragment Fab, Fab' or F(ab')<sub>2</sub>, a complementarity determining region, a soluble or membrane-anchored single-chain variable part, a single variable domain or a derivative of a monoclonal antibody according to any of claims 8 to 12 or of a humanized monoclonal antibody according to claim 13.
- 15 15. A pharmaceutical composition for the prevention or treatment of disorders of hemostasis and resulting pathologic conditions in mammals, comprising as an active ingredient a ligand according to any of claims 3 to 12 or a humanized monoclonal antibody according to claim 13 or an antigen-binding fragment Fab, Fab' or F(ab')<sub>2</sub>, or a soluble or membrane-anchored single-chain variable part or a single variable domain or a derivative according to claim 14, in admixture with a pharmaceutically acceptable carrier.
- 20 16. A pharmaceutical composition according to claim 15, wherein the only partial inactivation of a factor or a complex of two or more factors involved in hemostasis by the said active ingredient leads to a therapeutically useful plateau level of no more than 98% inactivation.
- 25 17. A pharmaceutical composition according to claim 15 or claim 16, wherein the only partial inactivation of a factor or a complex of two or more factors involved in hemostasis by the said active ingredient is at least 65% inactivation.
- 30 18. A pharmaceutical composition according to any of claims 15 to 17, further comprising a therapeutically effective amount of a thrombolytic agent.
- 35

19. A human hybrid or reshaped human monoclonal antibody which binds to and only partially inactivates factor VIII or a complex including factor VIII and which comprises only elements derived from the repertoire of human antibodies.
- 5 20. A human hybrid or reshaped human monoclonal antibody according to claim 19, wherein the said antibody interferes with a proteolytic cleavage site or the von Willebrand factor or the tenase complex reaction or by inducing a three-dimensional conformational change in factor VIII or by targeting a domain of factor VIII, in particular the C1 domain of factor VIII, or by targeting factor VIII in the factor VIII-  
10 von Willebrand factor complex.
21. A polynucleotide encoding for an antigen-binding fragment Fab, Fab' or F(ab')<sub>2</sub> or a complementarity determining region or a soluble or membrane-anchored single-chain variable part or a single variable domain or a derivative according to claim 14.
- 15 22. A method of obtaining monoclonal antibodies from a non-human mammal, comprising the steps of:
- a) selecting a non-human mammal having a modified and at least partially functional physiologically active protein, the modification being with respect to  
20 a wild type protein and lying in a domain of the protein;
  - b) administering the wild type protein to the non-human mammal in order to elicit an immune response, and
  - c) selecting B-lymphocytes from the non-human mammal which produce  
25 antibodies which only partially inactivate the wild type protein.
23. A method of obtaining monoclonal antibodies from the blood of a human being having a modified and at least partially functional physiologically active protein, the modification being with respect to a wild type protein and lying in a domain of the protein, and to whom the wild type protein was administered, the said method  
30 comprising the step of selecting, from the blood of said human being, B-lymphocytes which produce antibodies which only partially inactivate the wild type protein.
24. A method according to claim 22 or claim 23, wherein the wild type protein is a protein involved in a cascade of biochemical reactions.
- 35 25. A method according to any of claims 22 to 24, wherein the wild type protein is a factor or a complex of two or more factors involved in hemostasis.



26. A method according to any of claims 22 to 25, wherein the wild type protein is a factor or a complex of two or more factors involved in the coagulation cascade of blood.

5 27. A method according to any of claims 22 to 26, wherein the wild type protein is factor VIII or a complex including factor VIII.

28. A method according to any of claims 22 to 24, wherein the wild type protein is a protein involved in a proteolytic cascade.

10

29. A method according to any of claims 22, 23, 24 and 28, wherein the wild type protein is a complement factor.

30. A method of treatment and/or prevention of a disorder of hemostasis, coagulation disorder or thrombotic pathologic condition or attenuation of coagulation in a mammal, comprising administering to a mammal in need of such treatment or prevention or attenuation of coagulation a therapeutically effective amount of an active ingredient selected from a ligand according to any of claims 3 to 12, a humanized monoclonal antibody according to claim 13, an antigen-binding fragment Fab, Fab' or F(ab')<sub>2</sub>, a complementarity determining region, a soluble or membrane-anchored single-chain variable part, a single variable domain and a derivative according to claim 14.

15  
20

31. A method according to claim 30, wherein the thrombotic pathologic condition is selected from intravascular coagulation, arterial thrombosis, arterial restenosis, venous thrombosis and arteriosclerosis.

25

32. A method according to claim 30 or claim 31, wherein the active ingredient is provided to the mammal by oral, intranasal, subcutaneous, intramuscular, intradermal, intravenous, intraarterial or parenteral administration or by catheterization.

30

33. A method according to any of claims 30 to 32, further comprising administering to the mammal a therapeutically effective amount concomitantly or sequentially with a thrombolytic agent.

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
18 January 2001 (18.01.2001)

PCT

(10) International Publication Number  
**WO 01/04269 A1**

(51) International Patent Classification<sup>7</sup>: C12N 5/10, C07K  
16/36, 16/46, A61K 39/395, C12N 15/13, A61P 7/00

(74) Agents: BIRD, William, E. et al.; Bird Goën & Co, Vil-  
voordsebaan 92, B-3020 Winksele (BE).

(21) International Application Number: PCT/EP00/06677

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(22) International Filing Date: 13 July 2000 (13.07.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/143,891 14 July 1999 (14.07.1999) US  
9916450.1 14 July 1999 (14.07.1999) GB

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for all designated States except US*): LEUVEN  
RESEARCH & DEVELOPMENT VZW [BE/BE];  
Groot Begijnhof, Benedenstraat 59A, B-3000 Leuven  
(BE).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the  
claims and to be republished in the event of receipt of  
amendments.

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): JACQUEMIN,  
Marc, G. [BE/BE]; Rue Morimont 45, B-5330  
Sart-Bernard (BE). SAINT-REMY, Jean-Marie, R.  
[BE/BE]; Rue du Lambais 79, B-1390 Grez-Doiceau (BE).

*For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.*


(54) Title: LIGANDS FOR USE IN THERAPEUTIC COMPOSITIONS FOR THE TREATMENT OF HEMOSTASIS DISOR-  
DERS

(57) Abstract: The present invention comprises ligands and methods of manufacture thereof as well as pharmaceutical preparations including the ligands. The ligands may be human or humanized monoclonal antibodies and fragments, derivatives and homologs thereof. These may exhibit an unforeseen "plateau effect", i.e. the achievement of only a partial inactivation of a factor involved in hemostasis, in particular in the coagulation cascade, either individually or in combination even in molar excess. The ligands may bind to a factor or a complex of factors resulting in only partially impairing the function of a physiologically functional site of the said factor or factor complex even in molar excess. This makes the ligands particularly suitable for treating coagulation disorders and resulting thrombotic pathologic conditions while minimizing the risk of bleeding. Particularly useful is a property of ligands in accordance with the present invention to allow some physiological function of the affected site even when the ligand is in molar excess. The ligands may be anti-factor VIII antibodies or antibodies against a factor VIII complex, in particular human or human hybrid monoclonal antibodies which bind to factor VIII or a factor VIII complex and at least partially inhibit the activity of factor VIII.

WO 01/04269 A1

202050 22506001

IgG antibody binding to FVIII in ELISA



\* Inhibition of FVIII activity in coagulation assay

Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5

4/11

VH BO 2C11

1/1 31/11  
 atg gac tgg acc tgg agg atc ctc ttc ttg gtg gca gct aca ggc acc cac gcc cag  
 Met asp trp thr trp arg ile leu phe leu val ala ala thr gly thr his ala gln

61/21 91/31  
 gtc caa ctg gta cag tct ggg gct gag gtg aag cct ggg gcc tca gtg aag gtc tcc  
 val gln leu val gln ser gln ala glu val lys lys pro gly ala ser val lys val ser

121/41 151/51  
 tgc aag gtt tcc gga tac acc ctc act gaa tta ccc gtg cac tgg gtc gga cag gct cct  
 cys lys val ser gly tyr thr leu thr glu leu pro val his trp val gly gln ala pro  
 <-----CDR 1----->

181/61 211/71  
 gga aaa ggg ctt gag tgg gtg gga agt ttt gat cct gaa agt gga gaa tca atc tac gca  
 gly lys gly leu glu trp val gly ser phe asp pro glu ser gly glu ser ile tyr ala  
 <-----CDR 2----->

241/81 271/91  
 cgg gag ttc cag ggc agc gtc acc atg acc gcg gac aca tct aca gac ata gcc tac atg  
 arg glu phe gln gly ser val thr met thr ala asp thr ser thr asp ile ala tyr met  
 ----->

301/101 331/111  
 gag ctg agc agc ctg aga tct gac gac acg gcc gtg tat tac tgt gca gtc cct gac cct  
 glu leu ser ser leu arg ser asp asp thr ala val tyr tyr cys ala val pro asp pro  
 | <----->

361/121 391/131  
 gat gct ttt gat atc tgg ggc caa ggg aca atg gtc acc gtc tct tca gcc tcc acc aag  
 asp ala phe asp ile trp gly gln gly thr met val thr val ser ser ala ser thr lys  
 ---CDR 3----->

421/141  
 ggc cca tcg gtc ttc ccc ctg gga tcc cgt  
 gly pro ser val phe pro leu gly ser arg

Fig. 6

2025012250001

5/11

Fig. 7

VL BO 2C11

1/1

atg gaa acc cca gct cag ctt ctc ttc ctc ctg cta ctc tgg ctc cca gat acc acc gga  
 Met glu thr pro ala gln leu phe leu leu leu trp leu pro asp thr thr gly

31/11

61/21

gaa att gcg ttg acg cag tct cca ggc acc ctg tct ttg tct cca ggg gaa aga gcc acc  
 glu ile ala leu thr gln ser pro gly thr leu ser leu ser pro gly glu arg ala thr

91/31

121/41

ctc tcc tgc agg gcc agt cag agt ttt agc agc agc tac tta gcc tgg tat cag cag aaa  
 leu ser cys arg ala ser gln ser phe ser ser ser tyr leu ala trp tyr gln gln lys

151/51

181/61

cct ggc cag gct ccc agg ctc ctc atc tat ggt gca tcc acc agg gcc act ggc atc cca  
 pro gly gln ala pro arg leu leu ile tyr gly ala ser thr arg ala thr gly ile pro

211/71

241/81

gac agg ttc agt ggc agt ggg tct ggg aca gac ttc act ctc acc atc agc aga ctg gag  
 asp arg phe ser gly ser gly thr asp phe thr leu thr ile ser arg leu glu

271/91

301/101

cct gaa gat ttt gca gtg tat tac tgt cag aag tat ggt acg tca gcg atc acc ttc ggg  
 pro glu asp phe ala val tyr tyr cys gln lys tyr gly thr ser ala ile thr phe gly

331/111

361/121

caa ggg aca cga ctg gag att aaa gga act gtg gct gca cca tct gtc ttc atc ttc ccg  
 gln gly thr arg leu glu ile lys gly thr val ala ala pro ser val phe ile phe pro

391/131

421/141

cca tct  
 pro ser

202050 22505001

6/11

## VH Krlx-1 (Figure 8)

1/1  
 31/11  
 ATG GAC TGG ACC TGG AGG ATC CTC TTC TTG GTG GCA GCA GCC ACA GGA GCC CAC TCC CAG  
 Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Thr Gly Ala His Ser Gln

61/21  
 91/31  
 GTG CAA CTG GTG CAA TCT GGG GCT GAG GTG AAG AAG CCT GGG GCC TCA GTG AAG GTC TCC  
 Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys Pro Gly Ala Ser Val Lys Val Ser

121/41  
 151/51  
 TGC AAG ACC TCT GGA TAC AAC TTC ACC GGC TAC TCT GCT TCT GGA CAT ATC TTC ACC GCC  
 Cys Lys Thr Ser Gly Tyr Asn Phe Thr Gly Tyr Ser Ala Ser Gly His Ile Phe Thr Ala  
 <-----CDR1----->

181/61  
 211/71  
 TAC TCT GTG CAC TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT GAG TGG ATG GGA AGG ATC  
 Tyr Ser Val His Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met Gly Arg Ile  
 ----->

41/81  
 271/91  
 AAC CCT AAC AGT GGT GCC ACA GAC TAT GCA CAT AAA TTT CAG GCC AGG GTC ACC ATG TCC  
 Asn Pro Asn Ser Gly Ala Thr Asp Tyr Ala His Lys Phe Gln Gly Arg Val Thr Met Ser  
 -----CDR2----->

301/101  
 331/111



7/11

Fig. 8

AGG GAC ACG TCC ATC AGC ACA GCC TAC ATG GAA CTG AGC AGG CTG ACA TCT GAC GAC ACG  
 Arg Asp Thr Ser Ile Ser Thr Ala Tyr Met Glu Leu Ser Arg Leu Thr Ser Ser Asp Asp Thr

361/121

391/131

GCC ATG TAT TAC TGT GCG AGA GCC GAC AAC TAT TTC GAT ATT GTG ACT GGC TAT ACT TCT  
 Ala Met Tyr Tyr Cys Ala Arg Ala Asp Asn Tyr Phe Asp Ile Val Thr Gly Tyr Thr Ser

←-----CDR3----->

421/141

451/151

CAT TAC TTT GAC TAC TGG GCG CGG GGA ACC CTG GTC ACC GTC TCC TCA  
 His Tyr Phe Asp Tyr Trp Gly Arg Gly Thr Leu Val Thr Val Ser Ser

----->


8/11

VL KRIX 1


1/1  
 ATG GAA ACC CCA GCT CAG CTT CTC TTC CTC CTG CTA CTC TGG CTC CCA GAT ACC ACC GGA  
 Met glu thr pro ala gln leu leu phe leu leu leu trp leu pro asp thr thr gly  
 31/11  
 61/21  
 GAA ATT GTG TTG ACG CAG TTC CCA GGC ACC CTG TCT TTG TCT CCA GGG GAA AGA GCC ACC  
 glu ile val leu thr gln phe pro gly thr leu ser leu ser pro gly glu arg ala thr  
 91/31  
 121/41  
 CTC TCC TGC AGG GCC AGT CAG AGT GTT GCC AGC GCC TAC TTA GCC TGG TAC CAG CAA AAA  
 leu ser cys arg ala ser gln ser val ala ser ala tyr leu ala trp tyr gln gln lys  
 151/51  
 <-----CDR 1----->  
 181/61  
 CCT GGC CAG GCT CCC AGG CTC CTC ATC TAT GGT GCA TCC AGT AGG GCC ACC GAC ATC CCA  
 pro gly gln ala pro arg leu leu ile tyr gly ala ser ser arg ala thr asp ile pro  
 211/71  
 <-----CDR 2----->  
 241/81  
 CAC AGG TTC AGT GGC AGT GGC TCT GGG ACA GAC TTC ACT CTC ACC ATC AGC AGA CTG GAG  
 his arg phe ser gly ser gly ser gly thr asp phe thr leu thr ile ser arg leu glu  
 271/91  
 301/101  
 CCT GAA GAT TTT GCA GTG TAC TAC TGT CAG CAA TAT GGT ACC TCA GCC TTA CTC ACT TTC  
 pro glu asp phe ala val tyr tyr cys gln gln tyr gly thr ser ala leu leu thr phe  
 331/111  
 <-----CDR 3----->  
 361/121  
 GGC GGA GGC ACC AAG GTG GAG ATC AAA CGA ACT GTG GCT GCA CCA TCT GTC TTC ATC TTC  
 gly gly gly thr lys val glu ile lys arg thr val ala ala pro ser val phe ile phe  
 391/131  
 421/141  
 CCG CCA TCT  
 pro pro ser

Fig. 9


9/11

FIGURE 10

10/11

FIGURE 11

11/11  
FIGURE 12



WO 01/04269

PCT/EP00/06677

SEQUENCE LISTING

<110> LEUVEN RESEARCH & DEVELOPMENT vzw  
Jacquemin, Marc G  
Saint-Remy, Jean-Marie R

<120> LIGANDS FOR USE IN THERAPEUTIC COMPOSITIONS FOR THE  
TREATMENT OF HEMOSTASIS DISORDERS

<130> K1564-PCT

<140>

<141>

<150> GB9916450.1

<151> 1999-07-14

<150> US 60/143,891

<151> 1999-07-14

<160> 8

<170> PatentIn Ver. 2.1

<210> 1

<211> 143

<212> PRT

<213> human monoclonal antibody

<400> 1

Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Trp Leu Pro  
1 5 10 15

Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Phe Pro Gly Thr Leu Ser  
20 25 30

Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser  
35 40 45

Val Ala Ser Ala Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala  
50 55 60

Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Asp Ile Pro  
65 70 75 80

His Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile  
85 90 95

10030522-050202

Ser Arg Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr  
100 105 110

Gly Thr Ser Ala Leu Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile  
115 120 125

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser  
130 135 140

<210> 2

<211> 150

<212> PRT

<213> human monoclonal antibody

<400> 2

Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Ala Thr Gly  
1 5 10 15

Thr His Ala Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
20 25 30

Pro Gly Ala Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu  
35 40 45

Thr Glu Leu Pro Val His Trp Val Gly Gln Ala Pro Gly Lys Gly Leu  
50 55 60

Glu Trp Val Gly Ser Phe Asp Pro Glu Ser Gly Glu Ser Ile Tyr Ala  
65 70 75 80

Arg Glu Phe Gln Gly Ser Val Thr Met Thr Ala Asp Thr Ser Thr Asp  
85 90 95

Ile Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Asp Asp Thr Ala Val  
100 105 110

Tyr Tyr Cys Ala Val Pro Asp Pro Asp Ala Phe Asp Ile Trp Gly Gln  
115 120 125

Gly Thr Met Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val  
130 135 140

Phe Pro Leu Gly Ser Arg  
145 150

<210> 3

&lt;211&gt; 122

&lt;212&gt; PRT

&lt;213&gt; human monoclonal antibody

&lt;400&gt; 3

Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro  
 1 5 10 15

Asp Thr Thr Gly Glu Ile Ala Leu Thr Gln Ser Pro Gly Thr Leu Ser  
 20 25 30

Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser  
 35 40 45

Phe Ser Ser Ser Tyr Leu Ala Trp Tyr Gln Gln Lys Asp Arg Phe Ser  
 50 55 60

Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser Arg Leu Glu  
 65 70 75 80

Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Lys Tyr Gly Thr Ser Ala  
 85 90 95

Ile Thr Phe Gly Gln Gly Thr Arg Leu Glu Ile Lys Gly Thr Val Ala  
 100 105 110

Ala Pro Ser Val Phe Ile Phe Pro Pro Ser  
 115 120

&lt;210&gt; 4

&lt;211&gt; 429

&lt;212&gt; DNA

&lt;213&gt; human monoclonal antibody

&lt;220&gt;

&lt;221&gt; V\_region

&lt;222&gt; (1)..(429)

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (127)..(162)

&lt;223&gt; complementary determining region number one

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (205)..(225)

&lt;223&gt; complementary determining region number two



<220>  
 <221> misc\_feature  
 <222> (325)..(354)  
 <223> complementary determining region number three

<400> 4  
 atggaaaccc cagctcagct tctcttctc ctgctactct ggctcccaga taccaccgga 60  
 gaaatttgtg tgacgcagtt cccaggcacc ctgtctttgt ctccagggga aagagccacc 120  
 ctctcctgca gggccagtca gagtgttgcc agcgcctact tagcctggta ccagcaaaaa 180  
 cctggccagg ctcccaggct cctcatctat ggtgcatcca gtagggccac cgacatccca 240  
 cacaggttca gtggcagtgg gtctgggaca gacttcactc tcaccatcag cagactggag 300  
 cctgaagatt ttgcagtgtg ctactgtcag caatatggta cctcagcctt actcactttc 360  
 ggcggaggga ccaaggtgga gatcaaacga actgtggctg caccatctgt cttcatcttc 420  
 ccgccatct 429

<210> 5  
 <211> 450  
 <212> DNA  
 <213> human monoclonal antibody

<220>  
 <221> V\_region  
 <222> (1)..(450)

<220>  
 <221> misc\_feature  
 <222> (130)..(159)  
 <223> complementary determining region number one

<220>  
 <221> misc\_feature  
 <222> (202)..(258)  
 <223> complementary determining region number two

<220>  
 <221> misc\_feature  
 <222> (343)..(375)  
 <223> complementary determining region number three

<400> 5  
 atggactgga cctggaggat cctcttcttg gtggcagcag ctacaggcac ccacgcccag 60  
 gtccaactgg tacagtctgg ggctgaggtg aagaagcctg gggcctcagt gaaggctctcc 120  
 tgcaaggttt ccgatacac cctcactgaa ttaccctgac actgggtcgg acaggctctc 180  
 ggaaaagggc ttgagtgggt gggaagtgtt gatcctgaaa gtggagaatc aatctacgca 240  
 cgaggagttc agggcagcgt caccatgacc gcggacacat ctacagacat agcctacatg 300  
 gagctgagca gcctgagatc tgacgacacg gccgtgtatt actgtgcagt ccctgaccct 360

gatgcttttg atatctgggg ccaagggaca atggtcaccg tctcttcagc ctccaccaag 420  
ggcccatcgg tcttccccct gggatcccgt 450

<210> 6  
<211> 426  
<212> DNA  
<213> human monoclonal antibody

<220>  
<221> V\_region  
<222> (1)..(426)

<220>  
<221> misc\_feature  
<222> (7)..(162)  
<223> complementary determining region number one

<220>  
<221> misc\_feature  
<222> (205)..(225)  
<223> complementary determining region number two

<220>  
<221> misc\_feature  
<222> (325)..(351)  
<223> complementary determining region number three

<400> 6  
atggaaaccc cagctcagct tctcttcctc ctgctactct ggctcccaga taccaccgga 60  
gaaattgcgt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagagccacc 120  
ctctcctgca gggccagtc gagttttagc agcagctact tagcctggta tcagcagaaa 180  
cctggccagg ctcccaggct cctcatctat ggtgcatcca ccagggccac tggcatccca 240  
gacaggttca gtggcagtggt gtctgggaca gacttcactc tcaccatcag cagactggag 300  
cctgaagatt ttgcagtgtt ttactgtcag aagtatggta cgtcagcgat caccttcggg 360  
caagggacac gactggagat taaaggaaact gtggctgcac catctgtctt catcttccccg 420  
ccatct 426

<210> 7  
<211> 468  
<212> DNA  
<213> human monoclonal antibody

<220>  
<221> V\_region  
<222> (1)..(468)

<220>  
 <221> misc\_feature  
 <222> (124)..(192)  
 <223> complementary determining region number one

<220>  
 <221> misc\_feature  
 <222> (232)..(285)  
 <223> complementary determining region number two

<220>  
 <221> misc\_feature  
 <222> (282)..(435)  
 <223> complementary determining region number three

<400> 7  
 atggactgga cctggaggat cctcttcttg gtggcagcag ccacaggagc ccactcccag 60  
 gtgcaactgg tgcaatctgg ggctgaggtg aagaagcctg gggcctcagt gaaggtctcc 120  
 tgcaagacct ctggatacaa cttcaccggc tactctgctt ctggacatat cttcaccgcc 180  
 tactctgtgc actgggtgcg acaggcccct ggacaagggc ttgagtggat gggaaggatc 240  
 aaccctaaca gtggtgccac agactatgca cataaatttc agggcagggg caccatgtcc 300  
 agggacacgt ccatcagcac agcctacatg gaactgagca ggctgacatc tgacgacacg 360  
 gccatgtatt actgtgagag agccgacaac tatttcgata ttgtgactgg ctatacttct 420  
 cattactttg actactgggg ccggggaacc ctggtcaccg tctcctca 468

<210> 8  
 <211> 156  
 <212> PRT  
 <213> human monoclonal antibody

<400> 8  
 Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Ala Thr Gly  
 1 5 10 15  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 20 25 30  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Thr Ser Gly Tyr Asn Phe  
 35 40 45  
 Thr Gly Tyr Ser Ala Ser Gly His Ile Phe Thr Ala Tyr Ser Val His  
 50 55 60  
 Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met Gly Arg Ile  
 65 70 75 80  
 Asn Pro Asn Ser Gly Ala Thr Asp Tyr Ala His Lys Phe Gln Gly Arg

	85	90	95
Val Thr Met Ser Arg Asp Thr Ser Ile Ser Thr Ala Tyr Met Glu Leu			
	100	105	110
Ser Arg Leu Thr Ser Asp Asp Thr Ala Met Tyr Tyr Cys Ala Arg Ala			
	115	120	125
Asp Asn Tyr Phe Asp Ile Val Thr Gly Tyr Thr Ser His Tyr Phe Asp			
	130	135	140
Tyr Trp Gly Arg Gly Thr Leu Val Thr Val Ser Ser			
	145	150	155

0522-1769

**IN THE UNITED STATES PATENT AND TRADEMARK OFFICE**

IN RE THE APPLICATION OF )

Jaquemin, Marc G. & )  
Saint-Remy, Jean-Marie R. )

Examiner:

Group Art Unit:

SERIAL NO.: To Be Assigned )

FILED: )

FOR: "Ligands for Use in Therapeutic )  
Compositions for the Treatment of )  
Hemostasis Disorders" )

**CERTIFICATE OF MAILING**

I hereby certify that this correspondence is being deposited with the United States Postal Service as first class mail in an envelope addressed to " Director of Patents and Trademarks, Washington, D.C. 20231," on December 31, 2001.

Name of person signing Terri Dickinson

Signature

Honorable Director of  
Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

**PRIORITY CLAIM and ASSERTION OF SMALL ENTITY STATUS**

Dear Sir:

Applicants hereby claim priority benefits under 35 U.S.C. § 119 based upon Great Britain Application No. 9916450.1 and United States Provisional Application No. 60/143,891, both filed July 14, 1999. This application is the U.S. national phase application based upon International Application Serial No. PCT/EP00/06677, having international filing date of July 13, 2000, claiming priority upon said GB 9916450.1 and US 60/143,891.

Applicants hereby assert small entity status under 35 CFR § 1.27 with respect to the payment of fees for prosecution of the instant U.S. national phase application of International Application No. PCT/EP00/06677.

Respectfully Submitted,

December 31, 2001

Date



Mark A. Hagedorn  
Registration No. 44,731  
LEE, MANN, SMITH, MCWILLIAMS  
SWEENEY & OHLSON  
P.O. Box 2786  
Chicago, Illinois 60690-2786  
(312) 368-1300  
(312) 368-0034 (fax)

10030522, 060202

10/030522  
JC18 Rec'd PCT/PTO 31 DEC 2001

## SEQUENCE LISTING

<110> Leuven Research & Development vzw  
Jacquemin, Marc G  
Saint-Remy, Jean-Marie R

<120> Ligands for use in therapeutic compositions for the  
treatment of hemostatis disorders

<130> K1564-PCT

<150> GB9916450.1  
<151> 1999-07-14

<150> US 60/143,891  
<151> 1999-07-14

<160> 8

<170> PatentIn version 3.1

<210> 1  
<211> 143  
<212> PRT  
<213> Homo sapiens

<400> 1  
Met Glu Thr Pro Ala Gln Leu Leu Phe Leu Leu Leu Leu Trp Leu Pro  
1 5 10 15  
Asp Thr Thr Gly Glu Ile Val Leu Thr Gln Phe Pro Gly Thr Leu Ser  
20 25 30  
Leu Ser Pro Gly Glu Arg Ala Thr Leu Ser Cys Arg Ala Ser Gln Ser  
35 40 45  
Val Ala Ser Ala Tyr Leu Ala Trp Tyr Gln Gln Lys Pro Gly Gln Ala  
50 55 60  
Pro Arg Leu Leu Ile Tyr Gly Ala Ser Ser Arg Ala Thr Asp Ile Pro

65                                      70                                      75                                      80

His Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile

85                                      90                                      95

Ser Arg Leu Glu Pro Glu Asp Phe Ala Val Tyr Tyr Cys Gln Gln Tyr

100                                      105                                      110

Gly Thr Ser Ala Leu Leu Thr Phe Gly Gly Gly Thr Lys Val Glu Ile

115                                      120                                      125

Lys Arg Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser  
130                                      135                                      140

<210> 2  
<211> 150  
<212> PRT  
<213> Homo sapiens  
<400> 2

Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Ala Thr Gly

1                                      5                                      10                                      15

Thr His Ala Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys

20                                      25                                      30

Pro Gly Ala Ser Val Lys Val Ser Cys Lys Val Ser Gly Tyr Thr Leu

35                                      40                                      45

Thr Glu Leu Pro Val His Trp Val Gly Gln Ala Pro Gly Lys Gly Leu





	20		25		30										
Leu	Ser	Pro	Gly	Glu	Arg	Ala	Thr	Leu	Ser	Cys	Arg	Ala	Ser	Gln	Ser
	35		40		45										
Phe	Ser	Ser	Ser	Tyr	Leu	Ala	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Ala
50					55						60				
Pro	Arg	Leu	Leu	Ile	Tyr	Gly	Ala	Ser	Thr	Arg	Ala	Thr	Gly	Ile	Pro
65					70					75				80	
Asp	Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile
				85					90				95		
Ser	Arg	Leu	Glu	Pro	Glu	Asp	Phe	Ala	Val	Tyr	Tyr	Cys	Gln	Lys	Tyr
		100						105					110		
Gly	Thr	Ser	Ala	Ile	Thr	Phe	Gly	Gln	Gly	Thr	Arg	Leu	Glu	Ile	Lys
		115						120					125		
Gly	Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser		
130						135					140				

<210> 4  
 <211> 429  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> V\_region



&lt;223&gt;

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (130)..(159)

&lt;223&gt; complementary determining region number one

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (202)..(258)

&lt;223&gt; complementary determining region number two

&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (343)..(375)

&lt;223&gt; complementary determining region number three

&lt;400&gt; 5

atggactgga cctggaggat cctcttcttg gtggcagcag ctacaggcac ccacgccc  
ag 60gtccaactgg tacagtctgg ggctgagggt aagaagcctg ggcctcagt gaaggtct  
cc 120tgcaaggttt ccggatacac cctcactgaa ttacccgtgc actgggtcgg acaggctc  
ct 180ggaaaagggc ttgagtgggt ggaagtttt gatcctgaaa gtggagaatc aatctacg  
ca 240cgggagttcc agggcagcgt caccatgacc gcggacacat ctacagacat agcctaca  
tg 300gagctgagca gcctgagatc tgacgacacg gccgtgtatt actgtgcagt ccctgacc  
ct 360gatgcttttg atatctgggg ccaagggaca atggtcaccg tctcttcagc ctccacca  
ag 420ggcccatcgg tcttccccct gggatcccgt  
450

&lt;210&gt; 6

&lt;211&gt; 426

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; V\_region

&lt;222&gt; (1)..(426)

&lt;223&gt;

<220>  
 <221> misc\_feature  
 <222> (7)..(162)  
 <223> complementary determining region number one

<220>  
 <221> misc\_feature  
 <222> (205)..(225)  
 <223> complementary determining region number two

<220>  
 <221> misc\_feature  
 <222> (325)..(351)  
 <223> complementary determining region number three

<400> 6  
 atggaaaccc cagctcagct tctcttcctc ctgctactct ggctcccaga taccaccg  
 ga 60

gaaattgcgt tgacgcagtc tccaggcacc ctgtctttgt ctccagggga aagagcca  
 cc 120

ctctcctgca gggccagtca gagttttagc agcagctact tagcctggta tcagcaga  
 aa 180

cctggccagg ctcccaggct cctcatctat ggtgcatcca ccagggccac tggcatcc  
 ca 240

gacaggttca gtggcagtgg gtctgggaca gacttcactc tcaccatcag cagactgg  
 ag 300

cctgaagatt ttgcagtgta ttactgtcag aagtatggta cgtcagcgat caccttcg  
 gg 360

caagggacac gactggagat taaaggaact gtggctgcac catctgtctt catcttcc  
 cg 420

ccatct  
 426

<210> 7  
 <211> 468  
 <212> DNA  
 <213> Homo sapiens

<220>  
 <221> V\_region  
 <222> (1)..(468)  
 <223>

<220>  
 <221> misc\_feature  
 <222> (124)..(192)  
 <223> complementary determining region number one  
  
 <220>  
 <221> misc\_feature  
 <222> (232)..(285)  
 <223> complementary determining region number two  
  
 <220>  
 <221> misc\_feature  
 <222> (282)..(435)  
 <223> complementary determining region number three  
  
 <400> 7  
 atggactgga cctggaggat cctcttcttg gtggcagcag ccacaggagc ccactccc  
 ag 60  
  
 gtgcaactgg tgcaatctgg ggctgaggtg aagaagcctg gggcctcagt gaaggtct  
 cc 120  
  
 tgcaagacct ctggatacaa cttcaccggc tactctgctt ctggacatat cttcaccg  
 cc 180  
  
 tactctgtgc actgggtgcg acaggcccct ggacaagggc ttgagtggat gggaagga  
 tc 240  
  
 aaccctaaca gtggtgccac agactatgca cataaatttc agggcagggt caccatgt  
 cc 300  
  
 agggacacgt ccatcagcac agcctacatg gaactgagca ggctgacatc tgacgaca  
 cg 360  
  
 gccatgtatt actgtgcgag agccgacaac tatttcgata ttgtgactgg ctatactt  
 ct 420  
  
 cattactttg actactgggg ccggggaacc ctggtcaccg tctcctca  
 468  
  
 <210> 8  
 <211> 156  
 <212> PRT  
 <213> Homo sapiens  
  
 <400> 8  
 Met Asp Trp Thr Trp Arg Ile Leu Phe Leu Val Ala Ala Ala Thr Gly  
 1 5 10 15

Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 20 25 30

Pro Gly Ala Ser Val Lys Val Ser Cys Lys Thr Ser Gly Tyr Asn Phe  
 35 40 45

Thr Gly Tyr Ser Ala Ser Gly His Ile Phe Thr Ala Tyr Ser Val His  
 50 55 60

Trp Val Arg Gln Ala Pro Gly Gln Gly Leu Glu Trp Met Gly Arg Ile  
 65 70 75 80

Asn Pro Asn Ser Gly Ala Thr Asp Tyr Ala His Lys Phe Gln Gly Arg  
 85 90 95

Val Thr Met Ser Arg Asp Thr Ser Ile Ser Thr Ala Tyr Met Glu Leu  
 100 105 110

Ser Arg Leu Thr Ser Asp Asp Thr Ala Met Tyr Tyr Cys Ala Arg Ala  
 115 120 125

Asp Asn Tyr Phe Asp Ile Val Thr Gly Tyr Thr Ser His Tyr Phe Asp  
 130 135 140

Tyr Trp Gly Arg Gly Thr Leu Val Thr Val Ser Ser  
 145 150 155



02 MAY 2002

Docket No. 0522-1769**DECLARATION AND POWER OF ATTORNEY FOR PATENT APPLICATION**

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name.

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled LIGANDS FOR USE IN THERAPEUTIC COMPOSITIONS FOR THE TREATMENT OF HEMOSTASIS DISORDERS, the specification of which:

\_\_\_\_\_ is attached hereto.

X was filed on July 13, 2000 as Application Serial No. PCT/EP00/06677  
and was amended on \_\_\_\_\_ (if applicable).

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose information which is material to the examination of this application in accordance with Title 37, Code of Federal Regulations, Section 1.56(a).

I hereby claim priority benefits under Title 35, United States Code, Section 119 of any application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

**PRIOR APPLICATION(S)**

Country	Number	Date Filed	Priority Claimed	
			Yes	No
<u>GB</u>	<u>9916450.1</u>	<u>07/14/1999</u>	<u>x</u>	_____
<u>US</u>	<u>60/143,891</u>	<u>07/14/1999</u>	<u>x</u>	_____



[illegible]

And I hereby appoint Robert F. I. Conte, Registration No. 20,354, Thomas E. Smith, Registration No. 18,243, Dennis M. McWilliams, Registration No. 25,195, James R. Sweeney, Registration No. 18,721, William M. Lee, Jr., Registration No. 26,935, Glenn W. Ohlson, Registration No. 28,455, David C. Brezina, Registration No. 34,128, Jeffrey R. Gray, Registration No. 33,391, Timothy J. Engling, Registration No. 39,970, Gerald S. Geren, Registration No. 24,528, Peter J. Shakula, Registration No. 40,808, John W. Hayes, Registration No. 33,900, Wm. Marshall Lee, Registration No. 16,853, Mark A. Hagedorn, Registration No. 44,731, to prosecute this application and to transact all business in the Patent and Trademark Office connected herewith. It is requested that all communications be directed to:

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States

Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

1W Full name of sole or first inventor: Marc G. Jacquemin

Signature [Signature] Date 16.01.2002

Country of Residence: BELGIUM

Country of Citizenship: BELGIUM

Post Office and Residence Address: Rue Morimont 45

B-5330 Sart-Bernard

Belgium

BEY

2W Full name of joint inventor: Jean-Marie R. Saint-Remy

Signature [Signature] Date 16.1.2002

Country of Residence: BELGIUM

Country of Citizenship: BELGIUM

Post Office and Residence Address: Rue du Lambias 79

B-1390 Grez-Doiceau

Belgium

CBX